

**In-Common Laboratories**

Head Office: 57 Gervais Drive
North York, Ontario M3C 1Z2
(416) 422-3000
Toll Free: (888) 285-7817
www.ICLabs.ca

Bennett, Emily, ND
Bennett, Emily, ND
464 King St W
Hamilton, ON L8P 1B7
Canada

Patient Name

**Tigchelaar,
Alex**

Sex

M

Date of Birth (mm/dd/yyyy)

04/01/1982

Client's File No.

Order ID

56316279

Health Number

ICL Login Date (mm/dd/yyyy)

10/05/2016 12:49PM

Authorized Requester

Bennett, Emily, ND

Report Printed

11/11/2016 10:57AM

Comprehensive Stool Analysis with Parasitology x1

Sample ID: ICL161005059

Final - Approved 10/19/2016 2:25PM

Collection Date (mm/dd/yyyy)
10/03/2016 12:00PM

Order Choice Testing Site: Doctor's Data Inc., 3755 Illinois Avenue, St. Charles IL 60174

Comments:

TEST	RESULT	FLAG	NORMAL/THERAPEUTIC RANGE	UNITS	TEST SITE
Attached	Attachment				ICL1

Reporting Laboratories:

(1) ICL1, Non-Interfaced Test Site,

Patient Complete Name:
Tigchelaar, Alex

Order ID: 56316279

Current Page Number: 1

Total Pages Count: 1



LAB #: F161007-0103-1
PATIENT: Alex Tigchelaar
ID: P162810084
SEX: Male
DOB: 01/04/1982

AGE: 34

CLIENT #: 41676
DOCTOR: Emily Bennett, ND
In-Common Laboratories
57 Gervais Drive
North York, ON M3C 1Z2 CANADA

Comprehensive Stool Analysis / Parasitology x1

BACTERIOLOGY CULTURE

Expected/Beneficial flora

4+ Bacteroides fragilis group
2+ Bifidobacterium spp.
3+ Escherichia coli
2+ Lactobacillus spp.
3+ Enterococcus spp.

3+ Clostridium spp.
NG = No Growth

Commensal (Imbalanced) flora

4+ Alpha hemolytic strep
2+ Enterobacter cloacae complex
4+ Gamma hemolytic strep
2+ Klebsiella pneumoniae ssp pneumoniae

Dysbiotic flora

4+ Citrobacter freundii complex
4+ Citrobacter freundii complex, isolate 2

BACTERIA INFORMATION

Expected /Beneficial bacteria make up a significant portion of the total microflora in a healthy & balanced GI tract. These beneficial bacteria have many health-protecting effects in the GI tract including manufacturing vitamins, fermenting fibers, digesting proteins and carbohydrates, and propagating anti-tumor and anti-inflammatory factors.

Clostridia are prevalent flora in a healthy intestine. Clostridium spp. should be considered in the context of balance with other expected/beneficial flora. Absence of clostridia or over abundance relative to other expected/beneficial flora indicates bacterial imbalance. If *C. difficile* associated disease is suspected, a Comprehensive Clostridium culture or toxigenic *C. difficile* DNA test is recommended.

Commensal (Imbalanced) bacteria are usually neither pathogenic nor beneficial to the host GI tract. Imbalances can occur when there are insufficient levels of beneficial bacteria and increased levels of commensal bacteria. Certain commensal bacteria are reported as dysbiotic at higher levels.

Dysbiotic bacteria consist of known pathogenic bacteria and those that have the potential to cause disease in the GI tract. They can be present due to a number of factors including: consumption of contaminated water or food, exposure to chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

YEAST CULTURE

Normal flora

1+ Rhodotorula mucilaginosa

Dysbiotic flora

MICROSCOPIC YEAST

Result:

None

Expected:

None - Rare

The microscopic finding of yeast in the stool is helpful in identifying whether there is proliferation of yeast. Rare yeast may be normal; however, yeast observed in higher amounts (few, moderate, or many) is abnormal.

YEAST INFORMATION

Yeast normally can be found in small quantities in the skin, mouth, intestine and mucocutaneous junctions. Overgrowth of yeast can infect virtually every organ system, leading to an extensive array of clinical manifestations. Fungal diarrhea is associated with broad-spectrum antibiotics or alterations of the patient's immune status. Symptoms may include abdominal pain, cramping and irritation. When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast are not uniformly dispersed throughout the stool, this may lead to undetectable or low levels of yeast identified by microscopy, despite a cultured amount of yeast. Conversely, microscopic examination may reveal a significant amount of yeast present, but no yeast cultured. Yeast does not always survive transit through the intestines rendering it unviable.

Comments:

Date Collected: 10/03/2016

Date Received: 10/07/2016

Date Completed: 10/17/2016

* *Aeromonas, Campylobacter, Plesiomonas, Salmonella, Shigella, Vibrio, Yersinia, & Edwardsiella tarda* have been specifically tested for and found absent unless reported.





LAB #: F161007-0103-1
PATIENT: Alex Tigchelaar
ID: P162810084
SEX: Male
DOB: 01/04/1982

AGE: 34

CLIENT #: 41676
DOCTOR: Emily Bennett, ND
In-Common Laboratories
57 Gervais Drive
North York, ON M3C 1Z2 CANADA

Comprehensive Stool Analysis / Parasitology x1

PARASITOLOGY/MICROSCOPY

Sample 1

Rare Blastocystis hominis

PARASITOLOGY INFORMATION

Intestinal parasites are abnormal inhabitants of the gastrointestinal tract that have the potential to cause damage to their host. The presence of any parasite within the intestine generally confirms that the patient has acquired the organism through fecal-oral contamination. Damage to the host includes parasitic burden, migration, blockage and pressure. Immunologic inflammation, hypersensitivity reactions and cytotoxicity also play a large role in the morbidity of these diseases. The infective dose often relates to severity of the disease and repeat encounters can be additive.

There are two main classes of intestinal parasites, they include protozoa and helminths. The protozoa typically have two stages; the trophozoite stage that is the metabolically active, invasive stage and the cyst stage, which is the vegetative inactive form resistant to unfavorable environmental conditions outside the human host. Helminths are large, multicellular organisms. Like protozoa, helminths can be either free-living or parasitic in nature. In their adult form, helminths cannot multiply in humans.

In general, acute manifestations of parasitic infection may involve diarrhea with or without mucus and or blood, fever, nausea, or abdominal pain. However these symptoms do not always occur. Consequently, parasitic infections may not be diagnosed or eradicated. If left untreated, chronic parasitic infections can cause damage to the intestinal lining and can be an unsuspected cause of illness and fatigue. Chronic parasitic infections can also be associated with increased intestinal permeability, irritable bowel syndrome, irregular bowel movements, malabsorption, gastritis or indigestion, skin disorders, joint pain, allergic reactions, and decreased immune function.

In some instances, parasites may enter the circulation and travel to various organs causing severe organ diseases such as liver abscesses and cysticercosis. In addition, some larval migration can cause pneumonia and in rare cases hyper infection syndrome with large numbers of larvae being produced and found in every tissue of the body.

One negative parasitology x1 specimen does not rule out the possibility of parasitic disease, parasitology x3 is recommended. This test is not designed to detect *Cyclospora cayentanensis* or *Microsporidia* spp.

GIARDIA/CRYPTOSPORIDIUM IMMUNOASSAY

	Within	Outside	Reference Range
<i>Giardia duodenalis</i>	Neg		Neg
<i>Cryptosporidium</i>	Neg		Neg

Giardia duodenalis (AKA *intestinalis* and *lamblia*) is a protozoan that infects the small intestine and is passed in stool and spread by the fecal-oral route. Waterborne transmission is the major source of giardiasis.

Cryptosporidium is a coccidian protozoa that can be spread from direct person-to-person contact or waterborne transmission.

Comments:

Date Collected: 10/03/2016

Date Received: 10/07/2016

Date Reported: 10/17/2016



LAB #: F161007-0103-1
PATIENT: Alex Tigchelaar
ID: P162810084
SEX: Male
DOB: 01/04/1982

AGE: 34

CLIENT #: 41676
DOCTOR: Emily Bennett, ND
In-Common Laboratories
57 Gervais Drive
North York, ON M3C 1Z2 CANADA

Comprehensive Stool Analysis / Parasitology x1

DIGESTION / ABSORPTION

	Within	Outside	Reference Range
Elastase	> 500		> 200 µg/mL
Fat Stain	Few		None - Mod
Muscle fibers	None		None - Rare
Vegetable fibers	Rare		None - Few
Carbohydrates	Neg		Neg

Elastase findings can be used for the diagnosis or the exclusion of exocrine pancreatic insufficiency. Correlations between low levels and chronic pancreatitis and cancer have been reported. **Fat Stain:** Microscopic determination of fecal fat using Sudan IV staining is a qualitative procedure utilized to assess fat absorption and to detect steatorrhea. **Muscle fibers** in the stool are an indicator of incomplete digestion. Bloating, flatulence, feelings of "fullness" may be associated with increase in muscle fibers. **Vegetable fibers** in the stool may be indicative of inadequate chewing, or eating "on the run". **Carbohydrates:** The presence of reducing substances in stool specimens can indicate carbohydrate malabsorption.

INFLAMMATION

	Within	Outside	Reference Range
Lactoferrin	< 0.5		< 7.3 µg/mL
Calprotectin*	< 10		<= 50 µg/g
Lysozyme*	268		<= 600 ng/mL
White Blood Cells	None		None - Rare
Mucus	Neg		Neg

Lactoferrin and **Calprotectin** are reliable markers for differentiating organic inflammation (IBD) from functional symptoms (IBS) and for management of IBD. Monitoring levels of fecal lactoferrin and calprotectin can play an essential role in determining the effectiveness of therapy, are good predictors of IBD remission, and can indicate a low risk of relapse. **Lysozyme*** is an enzyme secreted at the site of inflammation in the GI tract and elevated levels have been identified in IBD patients. **White Blood Cells** (WBC) and **Mucus** in the stool can occur with bacterial and parasitic infections, with mucosal irritation, and inflammatory bowel diseases such as Crohn's disease or ulcerative colitis.

IMMUNOLOGY

	Within	Outside	Reference Range
Secretory IgA*		32.7	51 - 204 mg/dL

Secretory IgA* (sIgA) is secreted by mucosal tissue and represents the first line of defense of the GI mucosa and is central to the normal function of the GI tract as an immune barrier. Elevated levels of sIgA have been associated with an upregulated immune response.

Comments:

Date Collected: 10/03/2016
Date Received: 10/07/2016
Date Completed: 10/17/2016

*For Research Use Only. Not for use in diagnostic procedures.

Methodology: Elisa, Microscopy, Colormetric,
Gas Chromatography, pH Electrode



LAB #: F161007-0103-1
PATIENT: Alex Tigchelaar
ID: P162810084
SEX: Male
DOB: 01/04/1982

AGE: 34

CLIENT #: 41676
DOCTOR: Emily Bennett, ND
In-Common Laboratories
57 Gervais Drive
North York, ON M3C 1Z2 CANADA

Comprehensive Stool Analysis / Parasitology x1

SHORT CHAIN FATTY ACIDS

	Within	Outside	Reference Range
% Acetate	56		40 - 75 %
% Propionate	24		9 - 29 %
% Butyrate	16		9 - 37 %
% Valerate	3.3		0.5 - 7 %
Butyrate	2.3		0.8 - 4.8 mg/mL
Total SCFA's	14		4 - 18 mg/mL

Short chain fatty acids (SCFAs): SCFAs are the end product of the bacterial fermentation process of dietary fiber by beneficial flora in the gut and play an important role in the health of the GI as well as protecting against intestinal dysbiosis. Lactobacilli and bifidobacteria produce large amounts of short chain fatty acids, which decrease the pH of the intestines and therefore make the environment unsuitable for pathogens, including bacteria and yeast. Studies have shown that SCFAs have numerous implications in maintaining gut physiology. SCFAs decrease inflammation, stimulate healing, and contribute to normal cell metabolism and differentiation. Levels of **Butyrate** and **Total SCFA** in mg/mL are important for assessing overall SCFA production, and are reflective of beneficial flora levels and/or adequate fiber intake.

INTESTINAL HEALTH MARKERS

	Within	Outside	Reference Range
Red Blood Cells	None		None - Rare
pH	6.4		6 - 7.8
Occult Blood	Neg		Neg

Red Blood Cells (RBC) in the stool may be associated with a parasitic or bacterial infection, or an inflammatory bowel condition such as ulcerative colitis. Colorectal cancer, anal fistulas, and hemorrhoids should also be ruled out.

pH: Fecal pH is largely dependent on the fermentation of fiber by the beneficial flora of the gut.

Occult blood: A positive occult blood indicates the presence of free hemoglobin found in the stool, which is released when red blood cells are lysed.

MACROSCOPIC APPEARANCE

	Appearance	Expected
Color	Brown	Brown
Consistency	Soft	Formed/Soft

Color: Stool is normally brown because of pigments formed by bacteria acting on bile introduced into the digestive system from the liver. While certain conditions can cause changes in stool color, many changes are harmless and are caused by pigments in foods or dietary supplements. **Consistency:** Stool normally contains about 75% water and ideally should be formed and soft. Stool consistency can vary based upon transit time and water absorption.




LAB #: F161007-0103-1
PATIENT: Alex Tigchelaar
ID: P162810084
SEX: Male
AGE: 34

CLIENT #: 41676
DOCTOR: Emily Bennett, ND
In-Common Laboratories
57 Gervais Drive
North York, ON M3C 1Z2 CANADA

Bacterial Susceptibilities: *Citrobacter freundii* complex

NATURAL ANTIBACTERIALS



Natural antibacterial agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.

PRESCRIPTIVE AGENTS

	Resistant	Intermediate	Susceptible
Amoxicillin-Clavulanic Acid	R		
Ampicillin	R		
Cefazolin	R		
Ceftazidime			S
Ciprofloxacin			S
Trimeth-sulfa			S

Susceptible results imply that an infection due to the bacteria may be appropriately treated when the recommended dosage of the tested antimicrobial agent is used.

Intermediate results imply that response rates may be lower than for susceptible bacteria when the tested antimicrobial agent is used.

Resistant results imply that the bacteria will not be inhibited by normal dosage levels of the tested antimicrobial agent.

Comments:

Date Collected: 10/03/2016
Date Received: 10/07/2016
Date Completed: 10/17/2016

Natural antibacterial agent susceptibility testing is intended for research use only.
Not for use in diagnostic procedures.

v10.11




LAB #: F161007-0103-1
 PATIENT: Alex Tigchelaar
 ID: P162810084
 SEX: Male
 AGE: 34

CLIENT #: 41676
 DOCTOR: Emily Bennett, ND
 In-Common Laboratories
 57 Gervais Drive
 North York, ON M3C 1Z2 CANADA

Bacterial Susceptibilities: Citrobacter freundii complex, isolate 2

NATURAL ANTIBACTERIALS



Natural antibacterial agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.

PRESCRIPTIVE AGENTS

	Resistant	Intermediate	Susceptible
Amoxicillin-Clavulanic Acid	R		
Ampicillin	R		
Cefazolin	R		
Ceftazidime			S
Ciprofloxacin			S
Trimeth-sulfa			S

Susceptible results imply that an infection due to the bacteria may be appropriately treated when the recommended dosage of the tested antimicrobial agent is used.
Intermediate results imply that response rates may be lower than for susceptible bacteria when the tested antimicrobial agent is used.
Resistant results imply that the bacteria will not be inhibited by normal dosage levels of the tested antimicrobial agent.

Comments:

Date Collected: 10/03/2016
 Date Received: 10/07/2016
 Date Completed: 10/17/2016

Natural antibacterial agent susceptibility testing is intended for research use only.
 Not for use in diagnostic procedures.

v10.11




LAB #: F161007-0103-1
PATIENT: Alex Tigchelaar
ID: P162810084
SEX: Male
AGE: 34

CLIENT #: 41676
DOCTOR: Emily Bennett, ND
In-Common Laboratories
57 Gervais Drive
North York, ON M3C 1Z2 CANADA


Yeast Susceptibilities: *Rhodotorula mucilaginosa*

NATURAL ANTIFUNGALS



Natural antifungal agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.

NON-ABSORBED ANTIFUNGALS



Non-absorbed antifungals may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed using standardized commercially prepared disks impregnated with Nystatin. Relative sensitivity is reported based upon the diameter of the zone of inhibition surrounding the disk.

Comments:

Date Collected: 10/03/2016
Date Received: 10/07/2016
Date Completed: 10/17/2016

Yeast antifungal susceptibility testing is intended for research use only.
Not for use in diagnostic procedures.

v10.11

INTRODUCTION

This analysis of the stool specimen provides fundamental information about the overall gastrointestinal health of the patient. When abnormal microflora or significant aberrations in intestinal health markers are detected, specific interpretive paragraphs are presented. If no significant abnormalities are found, interpretive paragraphs are not presented.

Clostridium spp

Clostridia are expected inhabitants of the human intestine. Although most clostridia in the intestine are not virulent, certain species have been associated with disease. Clostridium perfringens is a major cause of food poisoning and is also one cause of antibiotic-associated diarrhea. Clostridium difficile is a causative agent in antibiotic-associated diarrhea and pseudomembranous colitis. Other species reported to be prevalent in high amounts in patients with Autistic Spectrum Disorder include Clostridium histolyticum group, Clostridium cluster I, Clostridium bolteae, and Clostridium tetani.

If these disease associations are a concern further testing may be necessary.

Washington W, Allen S, Janda W, Koneman E, Procop G, Schreckenberger P, Woods, G. Koneman's Color Atlas and Textbook of Diagnostic Microbiology, 6th edition. Lippincott Williams and Wilkins; 2006. pg 931-939

Song Y, Liu C, Finegold SM. Real-Time PCR Quantitation of Clostridia in Feces of Autistic Children. Applied and Environmental Microbiology. Nov. 2004, 6459-6465.

Parracho H, Bingham MO, Gibson GR, McCartney AL. Differences Between the Gut Microflora of Children with Autistic Spectrum Disorders and That of Healthy Children. Journal of Medical Microbiology. 2005;54, 987-991.

Imbalanced flora

Imbalanced flora are those bacteria that reside in the host gastrointestinal tract and neither injure nor benefit the host. Certain dysbiotic bacteria may appear under the imbalances category if found at low levels because they are not likely pathogenic at the levels detected. When imbalanced flora appear, it is not uncommon to find inadequate levels of one or more of the beneficial bacteria and/or a fecal pH which is more towards the alkaline end of the reference range (6 - 7.8). It is also not uncommon to find hemolytic or mucoid E. coli with a concomitant deficiency of beneficial E. coli and alkaline pH, secondary to a mutation of beneficial E. coli in alkaline conditions (DDI observations). Treatment with antimicrobial agents is unnecessary unless bacteria appear under the dysbiotic category.

Mackowiak PA. The normal microbial flora. N Engl J Med. 1982;307(2):83-93.

Dysbiotic Flora

In a healthy balanced state of intestinal flora, the beneficial bacteria make up a significant proportion of the total microflora. However, in many individuals there is an imbalance or deficiency of beneficial flora and an overgrowth of non-beneficial (imbalance) or even pathogenic microorganisms (dysbiosis). This can be due to a number of factors including: consumption of contaminated water or food; daily exposure of chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

A number of toxic substances can be produced by the dysbiotic bacteria including amines, ammonia, hydrogen sulfide, phenols, and secondary bile acids which may cause inflammation or damage to the brush border of the intestinal lining. If left unchecked, long-term damage to the intestinal lining may result in leaky gut syndrome, allergies, autoimmune disease (e.g. rheumatoid arthritis), irritable bowel syndrome, fatigue, chronic headaches, and sensitivities to a variety of foods. In addition, pathogenic bacteria can cause acute symptoms such as abdominal pain, nausea, diarrhea, vomiting, and fever in cases of food poisoning.

Bacterial sensitivities to a variety of prescriptive and natural agents have been provided for the pathogenic bacteria that were cultured from this patient's specimen. This provides the practitioner with useful information to help plan an appropriate treatment regimen. Supplementation with probiotics or consumption of foods (yogurt, kefir, miso, tempeh, tamari sauce) containing strains of lactobacilli, bifidobacteria, and enterococci can help restore healthy flora levels. Polyphenols in green and ginseng tea have been found to increase the numbers of beneficial bacteria. Hypochlorhydria may also predispose an individual to bacterial overgrowth, particularly in the small intestine. Nutritional anti-inflammatories can aid in reversing irritation to the GI lining. These include quercetin, vitamin C, curcumin, gamma-linoleic acid, omega-3 fatty acids (EPA, DHA), and aloe vera. Other nutrients such as zinc, beta-carotene, pantothenic acid, and L-glutamine provide support for regeneration of the GI mucosa. A comprehensive program may be helpful in individuals in whom a dysbiotic condition has caused extensive GI damage.

Lispi E. Digestive Wellness. New Canaan, CT: Keats Publishing; 1996.

Mitsuoka T. Intestinal Flora and Aging. Nutr Rev 1992;50(12):438-446.

Weisburger JH. Tea and Health: The Underlying Mechanisms. Proc Soc Exp Biol Med 1999;220(4):271-275.4.

Pereira SP, Gainsborough N, Dowling RH. Drug-induced Hypochlorhydria Causes High Duodenal Bacterial Counts in the Elderly. Ailment Pharmacol Ther 1998;12(1):99-104.

Murray MT. Stomach Ailments and Digestive Disturbances. Rocklin, CA: Prima Publishing; 1997.

Citrobacter species

Citrobacter species, a gram-negative bacterium and member of the Enterobacteriaceae family, is considered dysbiotic at 3+ or greater.

Citrobacter freundii complex, including Citrobacter freundii, Citrobacter braakii, Citrobacter gullenii, Citrobacter murlinae, Citrobacter rodentium, Citrobacter werkmanii, Citrobacter oungae, and less commonly, Citrobacter koseri and Citrobacter farmeri, can cause diarrheal disease. Symptoms due to Citrobacter freundii complex seem to be a result of the elaboration of an E. coli-like heat-stable enterotoxin and hydrogen sulfide. Citrobacter freundii complex has been implicated as a cause of gastrointestinal infection and inflammation, acute dysentery, and dyspepsia. Acute symptoms can include profuse, watery diarrhea which is often unaccompanied by abdominal pain, fecal blood, or white blood cells.

Citrobacter species thrive on Fructooligosaccharides (FOS), a common ingredient in artificial or alternative sweetener.

Antibiotics may be indicated if symptoms are prolonged. Refer to the bacterial sensitivities to identify the most appropriate agent.

Guarino, A, et al. Production of Escherichia coli Sta-Like Heat-Stable Enterotoxin by Citrobacter freundii Isolated from Humans. Journal of Clinical Microbiology. January 1987;110-114.

Washington W, Allen S, Janda W, Koneman E, Procop G, Schreckenberger P, Woods, G. Koneman's Color Atlas and Textbook of Diagnostic Microbiology, 6th edition. Lippincott Williams and Wilkins; 2006. pg 686-689.

Murray PR, Baron EJ, Jorgensen JH, Pfaller MA, Tenover FC, Tenover FC. Manual of Clinical Microbiology, 8th edition. Washington, DC: ASM Press; 2003. pg 701-704.

Cultured Yeast

Yeast, such as Candida are normally present in the GI tract in very small amounts. Many species of yeast exist and are commensal; however, they are always poised to create opportunistic infections and have detrimental effects throughout the body. Factors that contribute to a proliferation of yeast include frequent use of wide-spread antibiotics/low levels of beneficial flora, oral contraceptives, pregnancy, cortisone and other immunosuppressant drugs, weak immune system/low levels of sIgA, high-sugar diet, and high stress levels.

When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast grows in colonies and is typically not uniformly dispersed throughout the stool. This may lead to undetectable or low levels of yeast identified by microscopy, despite a cultured amount of yeast. Conversely, microscopic examination may reveal a significant amount of yeast present, but no yeast cultured. Yeast does not always survive transit through the intestines rendering it unviable for culturing. Therefore, both microscopic examination and culture are helpful in determining if abnormally high levels of yeast are present.

Parasites

Parasites were detected by microscopic examination in this stool specimen. Intestinal parasites are abnormal inhabitants of the GI tract that live off and have the potential to cause damage to their host. Factors such as contaminated food and water supplies, day care centers, increased international travel, pets, carriers such as mosquitoes and fleas, and sexual transmission have contributed to an increased prevalence of intestinal parasites.

In general, acute manifestations of parasitic infection may involve diarrhea with or without mucus and/or blood, fever, nausea, or abdominal pain. However, these symptoms do not always occur. Consequently, parasitic infections may not be diagnosed and eradicated. If left untreated, chronic parasitic infections can cause damage to the intestinal lining and can be an unsuspected cause of illness and fatigue. Chronic parasitic infections can also be associated with increased intestinal permeability, irritable bowel syndrome, irregular bowel movements, malabsorption, gastritis or indigestion, skin disorders, joint pain, allergic reactions, decreased immune function, and fatigue.

Murray MT. Stomach Ailments And Digestive Disturbances. Rocklin, CA: Prima Publishing;1997.

Gittleman AL. Guess What Came to Dinner Parasites And Your Health. New York, NY: Penguin Group; 2001.

Blastocystis hominis

Blastocystis hominis was identified in this specimen. Blastocystis hominis is a common protozoan found throughout the world. Blastocystis is transmitted via the fecal-oral route or from contaminated food or water.

Whether Blastocystis infection can cause symptoms is still considered controversial. Symptoms may be compounded by concomitant infection with other parasitic organisms, bacteria, or viruses. Often, B. hominis is found along with other such organisms. Nausea, diarrhea, abdominal pain, anal itching, weight loss, and excess gas have been reported in some persons with Blastocystis infection.

Metronidazole has been the traditionally considered the most effective drug (recommended adult dosage varies from 250 mg bid for 5-7 days to 750 mg tid x 10 days). Iodoquinol is also an effective medication (650 mg tid x 20 days). Recommended therapy can also eliminate G. lamblia, E. histolytica and D. fragilis. Various herbs may be effective, including oil of oregano. Limit refined carbohydrates in diet.

For more information:

1. Albrecht H, Stellbrink HJ, Koperski K, et al. Blastocystis hominis in human immunodeficiency virus-related diarrhea. Scand J Gastroenterol 1995;30:909-14.
2. Markell EK, Udkow MP. Blastocystis hominis: pathogen or fellow traveler Am J Trop Med Hyg 1986;35:1023-6.
3. Miller RA, Minshew BH. Blastocystis hominis: An organism in search of a disease. Rev Infect Dis 1988;10:930-8.

4. Udkow MP, Markell EK. Blastocystis hominis: prevalence in asymptomatic versus symptomatic hosts. *J Infect Dis* 1993;168:242-4.
5. Zuckerman MJ, Watts MT, Ho H., et al. Blastocystis hominis infection and intestinal injury. *Am J Med Sci* 1994;308:96-101.

References:

Sanford JP. The Sanford Guide to Antimicrobial Therapy. 35th edition. Gilbert DN, Moellering Jr, RC, Sande MA, eds. Hyde Park (VT): Antimicrobial Therapy Inc; 2005.

Abramowicz, M. The Medical Letter On Drugs and Therapeutics. Drugs For Parasitic Infections. New Rochelle (NY): The Medical Letter, Inc.

Beers, M. H., & Berkow, R. (Eds.). The Merck Manual of Diagnosis and Therapy Online. <http://www.merck.com/mrkshared/mmanual/section13/chapter161/161a.jsp>, Accessed August, 2005.

CDC Division of Parasitic Diseases website. <http://www.cdc.gov/ncidod/dpd/default.htm>, Accessed August, 2005.

Garcia, LS. Diagnostic Medical Parasitology. 4th ed. Washington DC: ASM; 2001; 6.

Leber AL, Movak SM In: Murray PR, Baron EJ, Pfaller MA, Tenover FC, Tenover FC, eds. Manual of Clinical Microbiology. 7th ed. Washington DC: ASM Press; 1999; 1401.

Secretory IgA (sIgA)

The concentration of sIgA is abnormally low in this specimen. Immunological activity in the gastrointestinal tract can be assessed using secretory immunoglobulin A (sIgA). Secretory IgA is the predominant antibody, or immune protein the body manufactures and releases in external secretions such as saliva, tears, and milk [1]. It is also transported through the epithelial cells that line the intestines out into the lumen. Secretory IgA represents the first line of defense of the GI mucosa and is central to the normal function of the GI tract as an immune barrier [1]. As the principal immunoglobulin isotype present in mucosal secretions, sIgA plays an important role in controlling intestinal milieu which is constantly presented with potentially harmful antigens such as pathogenic bacteria, parasites, yeast, viruses, abnormal cell antigens, and allergenic proteins [1]. Secretory IgA antibodies exert their function by binding to antigenic epitopes on the invading microorganism, limiting their mobility and adhesion to the epithelium of the mucus membrane [2]. This prevents the antigens from reaching systemic circulation and allowing them to be excreted directly in the feces.

Mental and physical stress as well as inadequate nutrition have been associated with low fecal sIgA concentrations. This includes dietary restrictions, excessive alcohol intake, body mass loss, negative moods, and anxiety [3]. One study found depressed levels of sIgA in malnourished children, particularly protein malnourishment, that responded well to nutritional rehabilitation with a significant increase in sIgA [4]. This may be because the synthesis and expression of

slgA requires adequate intake of the amino acid L-glutamine [3]. Animal studies have demonstrated that a glutamine-restricted diet can result in a 50% decrease in slgA levels [5]. An increase of dietary L-glutamine can restore GI immune function by protection of cells that synthesize slgA [6]. *Saccharomyces boulardii* is a nonpathogenic yeast that has been used for the treatment of acute infectious enteritis and antibiotic-associated diarrhea [7]. Significantly elevated levels of slgA and subsequent enhanced host immune response have been found following *S. boulardii* administration in mice and rats [8,9].

References:

1. Crago SS, Tomasi TB. Mucosal Antibodies, Food Allergy and Intolerance. Bailliere Tindall/W.B. Saunders 1987;167-89.
2. Roberts JA. Factors predisposing to urinary tract infections in children. *Ped Neph* 1996;10:517-522.
3. Carins J, Booth C. Salivary immunoglobulin-A as a marker of stress during strenuous physical training. *Aviat Space Environ Med* 2002;73(12)1203-7.
4. Teodosio MR, Oliveira ECM. Urinary secretory IgA after nutritional rehabilitation. *Braz J Med Biolog Res* 1999;32-421-426
5. Alverdy J. Effects of glutamine-supplemented diets on immunology of the gut. *J Parent Enteral Nutr* 1990;14(4):1095-1135.
6. Burke DJ, et al. Glutamine-supplemented total parenteral nutrition improves gut function. *Arch Surg* 1989;24:2396-2399.
7. Alverdy JA. The effect of total parenteral nutrition on gut lamina propria cells. *J Parent. Enteral Nutr* 1990;14(suppl).
8. Qamar A, Aboudola S, Warny M, et al. *Saccharomyces boulardii* stimulates intestinal immunoglobulin A immune response to clostridium difficile toxin A in mice. *Infect Immun* 2001;69(4):2762-5.
9. Buts JP, Bernasconi P, Vaerman JP, et al. Stimulation of secretory IgA and secretory component of immunoglobulins in small intestine of rats treated with *Saccharomyces boulardii*. *Dig Dis Sci* 1990;35(2):251-6.