



Metabolic Analysis Profile

Physician Copy



Parkgate House
356 West Barnes Lane
New Malden, Surrey KT3 6NB

63 Zillicoa Street
Asheville, NC 28801 USA



Results Overview

	Normal	Borderline	High Need	Supplementation for High Need
Antioxidants	<ul style="list-style-type: none"> Vitamin A / Carotenoids Vitamin C Vitamin E / Tocopherols 			
B-Vitamins	<ul style="list-style-type: none"> Riboflavin - B2 Niacin - B3 Biotin - B7 Folic Acid - B9 Cobalamin - B12 	<ul style="list-style-type: none"> Thiamin - B1 Pyridoxine - B6 		
Minerals	<ul style="list-style-type: none"> Magnesium Manganese Molybdenum Zinc 			

SUGGESTED SUPPLEMENT SCHEDULE

Supplements	Daily Recommended Intake (DRI)	Patient's Daily Recommendations	Provider Daily Recommendations
Antioxidants			
Vitamin A / Carotenoids	2,333 IU	3,000 IU	
Vitamin C	75 mg	250 mg	
Vitamin E / Tocopherols	22 IU	100 IU	
B-Vitamins			
Thiamin - B1	1.1 mg	25 mg	
Riboflavin - B2	1.1 mg	10 mg	
Niacin - B3	14 mg	20 mg	
Pyridoxine - B6	1.3 mg	25 mg	
Biotin - B7	30 mcg	100 mcg	
Folic Acid - B9	400 mcg	400 mcg	
Cobalamin - B12	2.4 mcg	100 mcg	
Minerals			
Magnesium	320 mg	400 mg	
Manganese	1.8 mg	3.0 mg	
Molybdenum	45 mcg	75 mcg	
Zinc	8 mg	10 mg	
Digestive Support			
Probiotics		10 B CFU	
Pancreatic Enzymes		0 IU	

Recommendations for age and gender-specific supplementation are set by comparing levels of nutrient functional need to optimal levels as described in the peer-reviewed literature. They are provided as guidance for short-term support of nutritional deficiencies only.

The Suggested Supplemental Schedule is provided at the request of the ordering practitioner. Any application of it as a therapeutic intervention is to be determined by the ordering practitioner.


Key

Normal	Borderline	High Need


Metabolic Analysis Profile Interpretation At-A-Glance

Nutritional Needs


Antioxidants



- ▶ Beta-carotene & other carotenoids are converted to vitamin A (retinol), involved in vision, antioxidant & immune function, gene expression & cell growth.
- ▶ Vitamin A deficiency may occur with chronic alcoholism, zinc deficiency, hypothyroidism, or oral contraceptives containing estrogen & progestin.
- ▶ Deficiency may result in night blindness, impaired immunity, healing & tissue regeneration, increased risk of infection, leukoplakia or keratosis.
- ▶ Food sources include cod liver oil, fortified cereals & milk, eggs, sweet potato, pumpkin, carrot, cantaloupe, mango, spinach, broccoli, kale & butternut squash.



- ▶ Vitamin C is an antioxidant (also used in the regeneration of other antioxidants). It is involved in cholesterol metabolism, the production & function of WBCs and antibodies, and the synthesis of collagen, norepinephrine and carnitine.
- ▶ Deficiency may occur with oral contraceptives, aspirin, diuretics or NSAIDs.
- ▶ Deficiency can result in scurvy, swollen gingiva, periodontal destruction, loose teeth, sore mouth, soft tissue ulcerations, or increased risk of infection.
- ▶ Food sources include oranges, grapefruit, strawberries, tomato, sweet red pepper, broccoli and potato.



- ▶ Alpha-tocopherol (body's main form of vitamin E) functions as an antioxidant, regulates cell signaling, influences immune function and inhibits coagulation.
- ▶ Deficiency may occur with malabsorption, cholestyramine, colestipol, isoniazid, orlistat, olestra and certain anti-convulsants (e.g., phenobarbital, phenytoin).
- ▶ Deficiency may result in peripheral neuropathy, ataxia, muscle weakness, retinopathy, and increased risk of CVD, prostate cancer and cataracts.
- ▶ Food sources include oils (olive, soy, corn, canola, safflower, sunflower), eggs, nuts, seeds, spinach, carrots, avocado, dark leafy greens and wheat germ.


Key

▶	Function
▶	Causes of Deficiency
▶	Complications of Deficiency
▶	Food Sources

Metabolic Analysis Profile Interpretation At-A-Glance

Nutritional Needs


B-Vitamins




- ▶ B1 is a required cofactor for enzymes involved in energy production from food, and for the synthesis of ATP, GTP, DNA, RNA and NADPH.
- ▶ Low B1 can result from chronic alcoholism, diuretics, digoxin, oral contraceptives and HRT, or large amounts of tea & coffee (contain anti-B1 factors).
- ▶ B1 deficiency may lead to dry beriberi (e.g., neuropathy, muscle weakness), wet beriberi (e.g., cardiac problems, edema), encephalopathy or dementia.
- ▶ Food sources include lentils, whole grains, wheat germ, Brazil nuts, peas, organ meats, brewer's yeast, blackstrap molasses, spinach, milk & eggs.



- ▶ B6 (as P5P) is a cofactor for enzymes involved in glycogenolysis & gluconeogenesis, and synthesis of neurotransmitters, heme, B3, RBCs and nucleic acids.
- ▶ Low B6 may result from chronic alcoholism, long-term diuretics, estrogens (oral contraceptives and HRT), anti-TB meds, penicillamine, L-DOPA or digoxin.
- ▶ B6 deficiency may result in neurologic symptoms (e.g., irritability, depression, seizures), oral inflammation, impaired immunity or increased homocysteine.
- ▶ Food sources include poultry, beef, beef liver, fish, whole grains, wheat germ, soybean, lentils, nuts & seeds, potato, spinach and carrots.




- ▶ B2 is a key component of enzymes involved in antioxidant function, energy production, detoxification, methionine metabolism and vitamin activation.
- ▶ Low B2 may result from chronic alcoholism, some anti-psychotic medications, oral contraceptives, tricyclic antidepressants, quinacrine or adriamycin.
- ▶ B2 deficiency may result in oxidative stress, mitochondrial dysfunction, low uric acid, low B3 or B6, high homocysteine, anemia or oral & throat inflammation.
- ▶ Food sources include milk, cheese, eggs, whole grains, beef, chicken, wheat germ, fish, broccoli, asparagus, spinach, mushrooms and almonds.




- ▶ Biotin is a cofactor for enzymes involved in functions such as fatty acid (FA) synthesis, mitochondrial FA oxidation, gluconeogenesis, and DNA replication & transcription.
- ▶ Deficiency may result from certain inborn errors, chronic intake of raw egg whites, long-term TPN use, anticonvulsants, high-dose B5, sulfa drugs & other antibiotics.
- ▶ Low levels may result in neurologic symptoms (e.g., paresthesias, depression), hair loss, scaly rash on face or genitals or impaired immunity.
- ▶ Food sources include yeast, whole grains, wheat germ, eggs, cheese, liver, meats, fish, wheat, nuts & seeds, avocado, raspberries, sweet potato and cauliflower.



- ▶ B3 is used to form NAD and NADP, involved in energy production from food, fatty acid & cholesterol synthesis, cell signaling, DNA repair & cell differentiation.
- ▶ Low B3 may result from deficiencies of tryptophan (B3 precursor), B6, B2 or Fe (cofactors in B3 production), or from long-term isoniazid or oral contraceptive use.
- ▶ B3 deficiency may result in pellagra (dermatitis, diarrhea, dementia), neurologic symptoms (e.g., depression, memory loss), bright red tongue or fatigue.
- ▶ Food sources include poultry, beef, organ meats, fish, whole grains, peanuts, seeds, lentils, brewer's yeast and lima beans.



- ▶ Folic acid plays a key role in coenzymes involved in DNA and SAMe synthesis, methylation, nucleic acids & amino acid metabolism and RBC production.
- ▶ Low folate may result from alcoholism, high-dose NSAIDs, diabetic meds, H2 blockers, some diuretics and anti-convulsants, SSRIs, methotrexate, trimethoprim, pyrimethamine, triamterene, sulfasalazine or cholestyramine.
- ▶ Folate deficiency can result in anemia, fatigue, low methionine, increased homocysteine, impaired immunity, heart disease, birth defects and CA risk.
- ▶ Food sources include fortified grains, green vegetables, beans & legumes.




- ▶ B12 plays important roles in energy production from fats & proteins, methylation, synthesis of hemoglobin & RBCs, and maintenance of nerve cells, DNA & RNA.
- ▶ Low B12 may result from alcoholism, malabsorption, hypochlorhydria (e.g., from atrophic gastritis, H. pylori infection, pernicious anemia, H2 blockers, PPIs), vegan diets, diabetic meds, cholestyramine, chloramphenicol, neomycin or colchicine.
- ▶ B12 deficiency can lead to anemia, fatigue, neurologic symptoms (e.g., paresthesias, memory loss, depression, dementia), methylation defects or chromosome breaks.
- ▶ Food sources include shellfish, red meat poultry, fish, eggs, milk and cheese.


Metabolic Analysis Profile Interpretation At-A-Glance

Nutritional Needs


Minerals



- ▶ Manganese plays an important role in antioxidant function, gluconeogenesis, the urea cycle, cartilage & bone formation, energy production and digestion.
- ▶ Impaired absorption of Mn may occur with excess intake of Fe, Ca, Cu, folic acid, or phosphorous compounds, or use of long-term TPN, Mg-containing antacids or laxatives.
- ▶ Deficiency may result in impaired bone/connective tissue growth, glucose & lipid dysregulation, infertility, oxidative stress, inflammation or hyperammonemia.
- ▶ Food sources include whole grains, legumes, dried fruits, nuts, dark green leafy vegetables, liver, kidney and tea.



- ▶ Magnesium is involved in >300 metabolic reactions. Key areas include energy production, bone & ATP formation, muscle & nerve conduction and cell signaling.
- ▶ Deficiency may occur with malabsorption, alcoholism, hyperparathyroidism, renal disorders (wasting), diabetes, diuretics, digoxin or high doses of zinc.
- ▶ Low Mg may result in muscle weakness/spasm, constipation, depression, hypertension, arrhythmias, hypocalcemia, hypokalemia or personality changes.
- ▶ Food sources include dark leafy greens, oatmeal, buckwheat, unpolished grains, chocolate, milk, nuts & seeds, lima beans and molasses.




- ▶ Molybdenum is a cofactor for enzymes that convert sulfites to sulfate, and nucleotides to uric acid, and that help metabolize aldehydes & other toxins.
- ▶ Low Mo levels may result from long-term TPN that does not include Mo.
- ▶ Mo deficiency may result in increased sulfite, decreased plasma uric acid (and antioxidant function), deficient sulfate, impaired sulfation (detoxification), neurologic disorders or brain damage (if severe deficiency).
- ▶ Food sources include buckwheat, beans, grains, nuts, beans, lentils, meats and vegetables (although Mo content of plants depends on soil content).




- ▶ Zinc plays a vital role in immunity, protein metabolism, heme synthesis, growth & development, reproduction, digestion and antioxidant function.
- ▶ Low levels may occur with malabsorption, alcoholism, chronic diarrhea, diabetes, excess Cu or Fe, diuretics, ACE inhibitors, H2 blockers or digoxin.
- ▶ Deficiency can result in hair loss and skin rashes, also impairments in growth & healing, immunity, sexual function, taste & smell and digestion.
- ▶ Food sources include oysters, organ meats, soybean, wheat germ, seeds, nuts, red meat, chicken, herring, milk, yeast, leafy and root vegetables.

Digestive Support




- ▶ Probiotics have many functions. These include: production of some B vitamins and vitamin K; enhancement of digestion & absorption; decreasing severity of diarrheal illness; modulation of immune function & intestinal permeability.
- ▶ Alterations of gastrointestinal microflora may result from C-section delivery, antibiotic use, improved sanitation, decreased consumption of fermented foods, and use of certain drugs.
- ▶ Some of the diseases associated with microflora imbalances include: IBS, IBD, fibromyalgia, chronic fatigue syndrome, obesity, atopic illness, colic and cancer.
- ▶ Food sources rich in probiotics are yogurt, kefir and fermented foods.



- ▶ Pancreatic enzymes are secreted by the exocrine glands of the pancreas and include protease/peptidase, lipase and amylase.
- ▶ Pancreatic exocrine insufficiency may be primary or secondary in nature. Any indication of insufficiency warrants further evaluation for underlying cause (i.e., celiac disease, small intestine villous atrophy, small bowel bacterial overgrowth).
- ▶ A high functional need for digestive enzymes suggests that there is an impairment related to digestive capacity.
- ▶ Determining the strength of the pancreatic enzyme support depends on the degree of functional impairment. Supplement potency is based on the lipase units present in both prescriptive and non-prescriptive agents.

Krebs Cycle At-A-Glance



All biomarkers reported in mmol/mol creatinine unless otherwise noted.

Metabolic Analysis Markers

Malabsorption and Dysbiosis Markers

Malabsorption Markers Reference Range

Indoleacetic Acid (IAA)	0.6	<= 4.2
Phenylacetic Acid (PAA)	0.04	<= 0.12

Bacterial Dysbiosis Markers

Dihydroxyphenylpropionic Acid (DHPPA)	3.2	<= 5.3
3-Hydroxyphenylacetic Acid	7.7	<= 8.1
4-Hydroxyphenylacetic Acid	6	<= 29
Benzoic Acid	0.04	<= 0.05
Hippuric Acid	83	<= 603

Yeast / Fungal Dysbiosis Markers

Arabinose	54	<= 96
Citramalic Acid	3.1	<= 5.8
Tartaric Acid	<dl	<= 15

Cellular Energy & Mitochondrial Metabolites

Carbohydrate Metabolism Reference Range

Lactic Acid	3.1	1.9-19.8
Pyruvic Acid	20	7-32
β-OH-Butyric Acid (BHBA)	14.4	<= 2.8

Energy Metabolism

Citric Acid	193	40-520
Cis-Aconitic Acid	15	10-36
Isocitric Acid	41	22-65
α-Ketoglutaric Acid (AKG)	26	4-52
Succinic Acid	1.4	0.4-4.6
Malic Acid	3.1	<= 3.0
β-OH-β-Methylglutaric Acid (HMG)	9	<= 15

Fatty Acid Metabolism

Adipic Acid	1.7	<= 2.8
Suberic Acid	0.6	<= 2.1

Creatinine Concentration

Reference Range

Creatinine ♦	13.5	3.1-19.5 mmol/L
--------------	------	-----------------

Neurotransmitter Metabolites

Reference Range

Vanilmandelic Acid	1.7	0.4-3.6
Homovanillic Acid	3.2	1.2-5.3
5-OH-indoleacetic Acid	8.4	3.8-12.1
3-Methyl-4-OH-phenylglycol	0.05	0.02-0.22
Kynurenic Acid	3.7	<= 7.1
Quinolinic Acid	5.2	<= 9.1
Kynurenic / Quinolinic Ratio	0.71	>= 0.44

Vitamin Markers

Reference Range

α-Ketoadipic Acid	0.6	<= 1.7
α-Ketoisovaleric Acid	0.63	<= 0.97
α-Ketoisocaproic Acid	0.55	<= 0.89
α-Keto-β-Methylvaleric Acid	0.8	<= 2.1
Formiminoglutamic Acid (FIGlu)	0.9	<= 1.5
Glutaric Acid	0.13	<= 0.51
Isovalerylglycine	2.0	<= 3.7
Methylmalonic Acid	0.8	<= 1.9
Xanthurenic Acid	0.81	<= 0.96
3-Hydroxypropionic Acid	6	5-22
3-Hydroxyisovaleric Acid	15	<= 29

Toxin & Detoxification Markers

Reference Range

α-Ketophenylacetic Acid (from Styrene)	0.18	<= 0.46
α-Hydroxyisobutyric Acid (from MTBE)	4.5	<= 6.7
Orotic Acid	0.49	0.33-1.01
Pyroglutamic Acid	22	16-34

Tyrosine Metabolism

Reference Range

Homogentisic Acid	12	<= 19
2-Hydroxyphenylacetic Acid	0.38	<= 0.76

Metabolic Analysis Reference Ranges are Age Specific

The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. Unless otherwise noted with ♦ as cleared by the U.S. Food and Drug Administration, assays are For Research Use Only.



Lipid Peroxides (Urine)



Parkgate House
356 West Barnes Lane
New Malden, Surrey KT3 6NB

63 Zillicoa Street
Asheville, NC 28801 USA



Lipid Peroxides (Urine)

Reference Range



Lab Comments

The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. Unless otherwise noted with ♦ as cleared by the U.S. Food and Drug Administration, assays are For Research Use Only.

Commentary is provided to the practitioner for educational purposes, and should not be interpreted as diagnostic or treatment recommendations. Diagnosis and treatment decisions are the responsibility of the practitioner.

Urine lipid peroxides is a marker of free radical damage in the body. An elevated level may reflect excess free radical production and/or insufficient antioxidants. Free radical damage is thought to underlie many processes such as atherosclerosis, chronic fatigue syndrome, cancer, cardiovascular disease, Parkinson's disease, Alzheimer's, and aging.