



# Comprehensive Digestive Stool Analysis



63 Zillicoa Street  
Asheville, NC 28801  
© Genova Diagnostics

Patient: **STEPHEN VAUGHAN**  
DOB: November 02, 1982  
Sex: M  
MRN: 1232709382

Order Number: **J9150230**  
Completed: July 27, 2016  
Received: July 15, 2016  
Collected: July 13, 2016

NutriPATH  
Mary Cavaggion  
18A Harker St  
Burwood  
Melbourne Victoria, 3125  
Australia

## Digestion

		Reference Range
Chymotrypsin		0.9-26.8 U/g
Putrefactive SCFAs (Total*)		1.3-8.6 micromol/g
* Total values equal the sum of all measurable parts.		
	Inside Outside	Reference Range
Meat Fibers		None
Vegetable Fibers		None - Few

## Absorption

		Reference Range
Triglycerides		0.2-3.3 mg/g
Long Chain Fatty Acids		1.3-23.7 mg/g
Cholesterol		0.2-3.5 mg/g
Phospholipids		0.2-8.8 mg/g
Fecal Fat (Total*)		2.6-32.4 mg/g
* Total values equal the sum of all measurable parts.		

## Metabolic Markers

Beneficial SCFAs (Total*)		Reference Range >= 13.6 micromol/g
n-Butyrate		= 2.5 micromol/g
Beta-Glucuronidase		337-4,433 U/g
pH		6.1-7.9

\* Total values equal the sum of all measurable parts.

## SCFA distribution

Acetate %		44.5-72.4 %
Propionate %		<= 32.1 %
n-Butyrate %		10.8-33.5 %

## Immunology

	Inside Outside	Reference Range
Fecal Lactoferrin ♦		Negative

## Macroscopic

Color		Brown
Mucus		Negative
Occult blood ♦		Negative

## Microbiology

### Bacteriology

#### Beneficial Bacteria

Lactobacillus species	
Escherichia coli	
Bifidobacterium	

#### Additional Bacteria

gamma haemolytic Streptococcus	NP	
alpha haemolytic Streptococcus	NP	
Haemolytic Escherichia coli	NP	

### Mycology

Candida albicans	NP	
------------------	----	--

\*NG



No Growth

NP



Non-Pathogen

PP



Possible Pathogen

P



Pathogen

Additional Tests (if indicated)

Commentary

Lab Comments  
SENSI'S: All yeast, add'l bacteria


The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. Unless otherwise noted with ♦, the assay has not been cleared by the U.S. Food and Drug Administration.

Commentary is provided to the practitioner for educational purposes, and should not be interpreted as diagnostic or treatment recommendations. Diagnosis and treatment decisions are the responsibility of the practitioner.

The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. Unless otherwise noted with ♦, the assay has not been cleared by the U.S. Food and Drug Administration.

The **Reference Range** is a statistical interval representing 95% or 2 Standard Deviations (2 S.D.) of the reference population.

One Standard Deviation (1 S.D.) is a statistical interval representing 68% of the reference population. Values between 1 and 2 S.D. are not necessarily abnormal. Clinical correlation is suggested. (See example below)



Human microflora is influenced by environmental factors and the competitive ecosystem of the organisms in the GI tract. Pathological significance should be based upon clinical symptoms and reproducibility of bacterial recovery.

Triglycerides constitute the major component of dietary fat and are normally broken down by pancreatic lipase into glycerol and free fatty acids. Triglycerides are within the reference range, indicating adequate fat digestion or a lack of dietary fat.

Chymotrypsin is within the reference range. Chymotrypsin is a key pancreatic enzyme that catalyzes protein digestion. Thus, the fecal level is a measure of proteolytic activity and a marker for pancreatic enzyme output as a whole. A value within the reference range suggests normal enzyme production. Levels are also influenced by transit time, such that faster transit results in higher fecal levels.

### Commentary

Valerate, iso-valerate and iso-butyrate are "putrefactive" short chain fatty acids, produced when anaerobic bacteria ferment undigested protein. Levels within the reference range suggest adequate protein digestion.

Long chain fatty acids (LCFAs) are within the reference range, suggesting adequate absorption of fats by the mucosa of the small intestine or a lack of dietary fat.

Cholesterol is within the reference range, suggesting adequate absorption of cholesterol by the small intestine or low dietary intake.

Phospholipids are normal. 50% of phospholipids are derived from bile, with 25% coming from mucosal desquamation and 25% from dietary sources. Nearly 85% of intestinal phospholipids are absorbed. Normal levels of fecal phospholipids indicate average dietary fat intake and adequate digestion/ absorption.

Total fecal fats are within the reference range. The total fecal fat is calculated as the sum of fecal triglycerides, phospholipids, cholesterol and long chain fatty acids.

Beneficial (Total) short chain fatty acids (SCFAs) are acetate, propionate and n-butyrate. They are the end products of anaerobic microbial fermentation of dietary fiber. Levels thus reflect the concentration of intestinal flora as well as soluble fiber in the diet. These beneficial SCFAs are crucial to the health of the intestine, serving as sources of fuel for the cells and the rest of the body. They also help to regulate the fluid balance in the colon.

n-Butyrate is the most important of the beneficial SCFAs, and is the primary energy source for colonic epithelial cells. Adequate amounts are necessary for the healthy metabolism of the colonic mucosa, and have been shown to have protective effects against colorectal cancers.

Beta-glucuronidase is within the reference range. This is an inducible enzyme, produced by E. coli and anaerobes Bacteroides, and Clostridia. Its activity reverses the detoxication of compounds processed in the hepatic Phase II glucuronidation pathway (including many pharmaceuticals, carcinogens, bile acids, and estrogen).

Fecal pH is within the reference range. The pH of the stool is a reflection of several factors in the GI tract, such as gastric acid, pancreatic bicarbonate, short chain fatty acids, ammonia, bile, organic acids, and acids produced by beneficial flora. Proper levels enhance colonization by beneficial flora, deter possible pathogens, promote normal digestive processes, and promote SCFA production.

The SCFA Distribution reflects the relative proportions of the beneficial SCFAs (n-butyrate, propionate, and acetate), thus providing an indirect measure of balance among the anaerobic organisms in the colon.

Sufficient amounts of Lactobacilli and E. coli appear to be present in the stool. Ample amounts of E. coli have been associated with a balanced gut flora. The "friendly bacteria", Lactobacilli and Bifidobacteria, are important for gastrointestinal function, as they are involved in vitamin synthesis, natural antibiotic production, immune defense, digestion, detoxification of pro-carcinogens and a host of other activities. Supplementation with Lactobacilli might be considered in selected cases where the organisms are in the low range of normal. Bifidobacteria is below optimal levels. Ideally, levels of Lactobacillus and E. coli should be 2+ or greater. Bifidobacteria being a predominate anaerobe should be recovered at levels of 4+.

There is no detection of fecal lactoferrin. This indicates no active intestinal inflammation. However, non-inflammatory diarrhea caused by irritable bowel syndrome, small intestinal viral infections, non-invasive parasitic infections, or other etiologies may still be present even in the absence of lactoferrin.



Patient: **STEPHEN  
VAUGHAN**  
DOB: November 02, 1982  
Sex: M  
MRN: 1232709382

Order Number: **J9150230**

Completed: July 27, 2016

Received: July 15, 2016

Collected: July 13, 2016



63 Zillico Street  
Asheville, NC 28801  
© Genova Diagnostics

## Azole Antifungals

### CANDIDA ALBICANS

	R	I	S-DD*	S	NI*
Fluconazole	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text" value="≤1"/>	<input type="text"/>
Voriconazole	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text" value="≤0.12"/>	<input type="text"/>

## Non-absorbed Antifungals

### CANDIDA ALBICANS

	Low Inhibition	High Inhibition
Nystatin	<input type="text"/>	<input type="text"/>

## Natural Antifungals

### CANDIDA ALBICANS

	Low Inhibition	High Inhibition
Berberine	<input type="text"/>	<input type="text"/>
Caprylic Acid	<input type="text"/>	<input type="text"/>
Garlic	<input type="text"/>	<input type="text"/>
Undecylenic Acid	<input type="text"/>	<input type="text"/>
Plant tannins	<input type="text"/>	<input type="text"/>
Uva-Ursi	<input type="text"/>	<input type="text"/>

#### Prescriptive Agents:

The R (Resistant) category implies isolate is not inhibited by obtainable levels of pharmaceutical agent.

The I (Intermediate) category includes isolates for which the minimum inhibition concentration (MIC) values usually approach obtainable pharmaceutical agent levels and for which response rates may be lower than for susceptible isolates.

\* The S-DD (Susceptible-Dose Dependent) category implies clinical efficacy when higher than normal dosage of a drug can be used and maximal concentration achieved.

The S (Susceptible) column implies that isolates are inhibited by the usually achievable concentrations of the pharmaceutical agent.

\* NI (No Interpretive guidelines established) category is used for organisms that currently do not have established guidelines for MIC interpretation.

Refer to published pharmaceutical guidelines for appropriate dosage therapy.

#### Nystatin and Natural Agents:

Results for Nystatin are being reported with natural antifungals in this category in accordance with laboratory guidelines for reporting sensitivities. In this assay, inhibition is defined as the reduction level on organism growth as a direct result of inhibition by a natural substance. The level of inhibition is an indicator of how effective the substance was at limiting the growth of an organism in an in vitro environment. High inhibition indicates a greater ability by the substance to limit growth, while Low Inhibition a lesser ability to limit growth. The designated natural products should be considered investigational in nature and not be viewed as standard clinical treatment substances.

This test has been developed and its performance characteristics determined by Genova Diagnostics, Inc. It has not been cleared or approved by the U.S. Food and Drug Administration.