

# livewello

## Gene Variance report

Uri rahamim / F

### ALLERGY

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	C11orf30	rs2155219	G	TT	-/-
2	HLA-DQB1	rs7775228	C	TT	-/-

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### CLEFT LIP/CLEFT PALATE

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	IRF6	rs861020	A	GG	-/-
2	PVT1	rs987525	A	CC	-/-
3	RARA	rs7217852	G	AG	+/-
4	RARA	rs9904270	T	—	NG
5	TBX22	rs41307258	-	TT	

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### CLOTTING FACTORS

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	CETP	rs1800775	C	AC	+/-
2	CYP4V2	rs13146272	C	AC	+/-
3	F10	rs3211719	G	AA	-/-
4	F11	rs2036914	T	—	NG
5	F11	rs2289252	T	CC	-/-
6	F12	rs1801020	A	—	NG
7	F12	rs2731672	T	CC	-/-
8	F3	rs1324214	A	—	NG
9	F5	rs6025	T	CC	-/-
11	F9	rs6048	-	AA	
12	GP6	rs1613662	G	AG	+/-
13	HRG	rs9898	T	CT	+/-
14	ITGB3	rs5918	C	CT	+/-
15	NR1I2	rs1523127	C	AC	+/-
16	SERPINC1	rs2227589	T	CC	-/-

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### DETOX

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	CTH	rs1021737	T	GT	+/-
3	CYP1A1	rs1799814	T	GG	-/-
4	CYP1A1	rs4986883	C	TT	-/-
5	CYP1A2	rs762551	C	AA	-/-
8	CYP1B1	rs1800440	C	CT	+/-
11	CYP2C9	rs1057910	C	AA	-/-
12	CYP2C9	rs1799853	T	CT	+/-
13	CYP2D6	rs1065852	A	GG	-/-
15	CYP2D6	rs16947	A	AG	+/-
16	CYP2E1	rs2070676	G	CC	-/-
18	CYP2E1	rs6413419	A	GG	-/-
20	CYP3A4	rs2740574	C	TT	-/-
21	CYP3A4	rs4986910	G	AA	-/-
22	CYP3A4	rs55785340	G	AA	-/-
23	GPX3	rs8177412	C	—	NG
24	GSTM1	rs1056806	T	CC	-/-
25	GSTM1	rs2239892	G	—	NG
26	GSTM1	rs4147565	A	--	NC
27	GSTM1	rs4147567	G	—	NG
28	GSTP1	rs1138272	T	CC	-/-
29	GSTP1	rs1695	G	AA	-/-
30	NAT1	rs4986782	A	GG	-/-
31	NAT2	rs1208	G	AA	-/-
32	NAT2	rs1799930	A	AG	+/-
33	NAT2	rs1799931	A	AG	+/-
34	NAT2	rs1801279	A	GG	-/-
35	NAT2	rs1801280	C	TT	-/-

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified

health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

# Uri rahamim / F

## GLUTEN INTOLERANCE

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	HLA-DQA1	rs2187668	T	CT	+/-
2	HLA-DQA2	rs2858331	G	AA	-/-

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### IGA

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	CFH	rs6677604	A	GG	-/-
2	HLA-DPA1	rs1883414	A	AG	+/-
3	HLA-DQA2	rs9275224	A	AA	+/+
4	HLA-DRB1	rs9275596	C	CT	+/-
5	HORMAD2	rs2412971	G	AA	-/-
6	IFIH1	rs1990760	T	TT	+/+
7	IGF1R	rs2229765	A	GG	-/-
8	IRF5	rs4728142	A	AG	+/-
9	MHC	rs9271366	G	AA	-/-
10	MTC03P1	rs2856717	A	AA	+/+
11	PSMB8	rs9357155	A	AG	+/-
12	TRAF1	rs3761847	G	AA	-/-

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### IGE

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	C3	rs10402876	C	CG	+/-
3	CD14	rs2569191	C	TT	-/-
4	DARC	rs2814778	C	TT	-/-
5	FCER1A	rs2251746	C	CC	+/+
6	FCER1A	rs2427824	T	CC	-/-
7	FCER1A	rs2427827	-	CC	
8	FCER1A	rs2427837	A	—	NG
9	FCER1A	rs2494262	A	—	NG
10	FCER1A / OR10J2P	rs10489854	T	CC	-/-
11	IL13	rs1295685	A	—	NG
12	IL13	rs1800925	T	CC	-/-
13	IL5	rs2069812	G	AG	+/-
14	RAD50	rs17772565	T	CC	-/-
15	RAD50	rs17772583	G	AA	-/-
16	RAD50	rs2040704	G	AA	-/-
17	RAD50	rs2240032	T	CC	-/-
18	RAD50	rs6884762	T	CC	-/-
19	RAG1	rs3740955	A	AA	+/+
20	SOCS1	rs33977706	A	AC	+/-

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### IGG

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	FCGR2A	rs1801274	G	AG	+/-
2	GSTM3	rs7483	T	CT	+/-
3	Intergenic	rs2013111	C	—	NG
4	MUC21	rs1634731	G	AA	-/-
5	TBC1D27	rs3751987	A	—	NG
6	TNFRSF13B	rs4792800	G	—	NG

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### METHYLATION

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	ACAT1	rs3741049	A	GG	-/-
2	ACE	rs4343	G	GG	+/+
3	AGT	rs699	A	AA	+/+
4	AHCY-01	rs819147	C	TT	-/-
5	AHCY-02	rs819134	G	—	NG
6	AHCY-19	rs819171	C	TT	-/-
7	BHMT	rs3733890	A	AG	+/-
8	BHMT-02	rs567754	T	CC	-/-
9	BHMT-04	rs617219	C	—	NG
10	BHMT-08	rs651852	T	CC	-/-
11	C1orf167	rs4846048	G	AG	+/-
12	CBS	rs2851391	T	CC	-/-
13	CBS	rs4920037	A	GG	-/-
14	CBS A360A	rs1801181	A	AG	+/-
15	CBS C699T	rs234706	A	GG	-/-
16	CBS N212N	rs2298758	A	—	NG
17	CLCN6	rs13306560	T	CC	-/-
18	CLCN6	rs13306561	G	—	NG
20	COMT	rs6269	G	AG	+/-
21	COMT H62H	rs4633	T	CT	+/-
22	COMT P199P	rs769224	A	GG	-/-
23	COMT V158M	rs4680	A	AG	+/-
24	DAO	rs2070586	A	GG	-/-
25	DAO	rs2111902	G	—	NG
26	DAO	rs3741775	C	AC	+/-
27	DHFR	rs1643649	C	CC	+/+
28	FOLR1	rs2071010	A	GG	-/-
29	FOLR2	rs651933	G	AA	-/-
30	FOLR3	rs7925545	G	AA	-/-
31	FOLR3	rs7926875	A	—	NG
32	FUT2	rs492602	G	AG	+/-
33	FUT2	rs601338	A	AG	+/-
34	FUT2	rs602662	A	AG	+/-

	Gene	rsID	Minor Allele	Genotype	Phenotype
35	G6PD	rs1050828	-	CC	
36	G6PD	rs1050829	-	TT	
37	GAD1	rs10432420	A	AG	+/-
38	GAD1	rs12185692	A	CC	-/-
39	GAD1	rs2058725	C	TT	-/-
40	GAD1	rs2241165	C	CT	+/-
41	GAD1	rs3749034	A	AG	+/-
42	GAD1	rs3791850	A	GG	-/-
43	GAD1	rs3791851	C	CT	+/-
44	GAD1	rs3791878	T	GT	+/-
45	GAD1	rs3828275	T	CC	-/-
46	GAD1	rs701492	T	CT	+/-
47	GAD1	rs769395	G	—	NG
48	GAD1	rs769407	C	CG	+/-
49	GAD2	rs1805398	T	—	NG
50	GAMT	rs17851582	A	GG	-/-
51	GAMT	rs55776826	T	CC	-/-
52	GIF	rs558660	A	AG	+/-
53	MAOA	rs6323	-	TT	
54	MAOB	rs1799836	-	TT	
55	MTHFD1	rs1076991	T	CT	+/-
56	MTHFD1	rs2236225	A	GG	-/-
57	MTHFD1L	rs11754661	A	GG	-/-
58	MTHFD1L	rs17349743	C	CT	+/-
59	MTHFD1L	rs6922269	A	GG	-/-
60	MTHFD1L	rs803422	A	GG	-/-
61	MTHFR	rs12121543	-	—	
62	MTHFR	rs1476413	T	CC	-/-
63	MTHFR	rs17037390	A	GG	-/-
64	MTHFR	rs17037396	T	—	NG
65	MTHFR	rs17367504	G	AA	-/-
66	MTHFR	rs2066470	A	GG	-/-
67	MTHFR	rs2274976	T	CC	-/-
68	MTHFR	rs4846049	T	GT	+/-
69	MTHFR A1298C	rs1801131	G	TT	-/-
70	MTHFR C677T	rs1801133	A	GG	-/-
71	MTHFS	rs6495446	T	CT	+/-

	Gene	rsID	Minor Allele	Genotype	Phenotype
72	MTR A2756G	rs1805087	G	AG	+/-
73	MTRR	rs10380	T	—	NG
74	MTRR	rs1801394	G	AA	-/-
75	MTRR A664A	rs1802059	A	AG	+/-
76	MTRR K350A	rs162036	G	AG	+/-
77	MTRR R415T	rs2287780	T	—	NG
78	NOS1	rs3782206	T	—	NG
79	NOS2	rs2248814	A	AG	+/-
80	NOS2	rs2274894	T	GT	+/-
81	NOS2	rs2297518	A	GG	-/-
82	NOS3	rs1800779	G	AG	+/-
83	NOS3	rs1800783	A	AT	+/-
84	NOS3	rs2070744	C	—	NG
86	NOS3	rs7830	T	—	NG
87	PEMT	rs4244593	T	TT	+/+
89	SHMT1	rs1979277	A	—	NG
90	SLC19A1	rs1888530	C	—	NG
91	SLC19A1	rs3788200	A	—	NG
92	SOD2	rs2758331	A	AA	+/+
93	SOD2	rs4880	G	GG	+/+
94	SOD3	rs2855262	C	CT	+/-
95	TCN1	rs526934	G	AG	+/-
96	TCN2	rs1801198	G	CG	+/-
97	TYMSOS	rs502396	T	—	NG
98	VDR Bsm	rs1544410	T	CC	-/-
99	VDR Taq	rs731236	G	AA	-/-

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### MITOCHONDRIAL FUNCTION

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	ATP5C1	rs1244414	T	CC	-/-
2	ATP5C1	rs1244422	T	—	NG
3	ATP5C1	rs12770829	T	—	NG
4	ATP5C1	rs2778475	A	—	NG
5	ATP5C1	rs4655	C	—	NG
6	ATP5G3	rs185584	G	—	NG
7	ATP5G3	rs36089250	C	TT	-/-
8	CCL2	rs1024611	G	AA	-/-
9	COX5A	rs8042694	G	AA	-/-
10	COX6C	rs1135382	A	—	NG
11	COX6C	rs12544943	G	—	NG
12	COX6C	rs4510829	A	—	NG
13	COX6C	rs4626565	C	TT	-/-
14	COX6C	rs7828241	C	—	NG
15	COX6C	rs7844439	A	—	NG
16	NDUFS3	rs2233354	C	—	NG
17	NDUFS3	rs4147730	A	GG	-/-
18	NDUFS3	rs4147731	A	—	NG
19	NDUFS7	rs1142530	T	CT	+/-
20	NDUFS7	rs11666067	A	—	NG
21	NDUFS7	rs2074895	A	—	NG
22	NDUFS7	rs2332496	A	AG	+/-
23	NDUFS7	rs7254913	G	—	NG
24	NDUFS7	rs7258846	T	GT	+/-
25	NDUFS7	rs809359	G	AA	-/-
27	NDUFS8	rs1104739	C	—	NG
28	NDUFS8	rs1122731	A	—	NG
29	NDUFS8	rs2075626	C	CC	+/+
30	NDUFS8	rs3115546	G	—	NG
31	NDUFS8	rs4147776	C	—	NG
32	NDUFS8	rs999571	A	GG	-/-
33	SLC19A1	rs1051266	C	TT	-/-
34	UQCRC2	rs11648723	T	GG	-/-

	Gene	rsID	Minor Allele	Genotype	Phenotype
35	UQCRC2	rs4850	A	GG	-/-
36	UQCRC2	rs6497563	C	—	NG

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### OTHER IMMUNE FACTORS

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	ADD1	rs4961	T	GG	-/-
2	APOE	rs429358	C	TT	-/-
3	ATG16L1	rs10210302	T	TT	+/+
4	HLA-DRB1	rs660895	G	AA	-/-
5	IL13	rs20541	A	AG	+/-
6	IL4R	rs1801275	G	AG	+/-
7	KIAA1109	rs6822844	T	GG	-/-
8	MEFV	rs11466023	A	GG	-/-
10	STAT4	rs10181656	G	CC	-/-
11	TNF	rs1800629	A	GG	-/-
12	TNF	rs361525	A	GG	-/-

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

## Uri rahamim / F

### SULFOTRANSFERASE

	Gene	rsID	Minor Allele	Genotype	Phenotype
2	SULT1A1	rs1801030	-	—	
3	SULT1A1	rs4149381	G	TT	-/-
4	SULT1A1	rs6498090	A	—	NG
5	SULT1A1	rs7192559	T	CC	-/-
6	SULT1A1	rs7193599	C	—	NG
7	SULT1A1	rs8057055	A	—	NG
8	SULT1A1	rs9282862	C	TT	-/-
9	SULT2A1	rs11083907	A	GG	-/-
10	SULT2A1	rs11569679	T	CC	-/-
11	SULT2A1	rs2547231	C	AA	-/-
12	SULT2A1	rs2547242	C	—	NG
13	SULT2A1	rs2910393	T	—	NG
15	SULT2A1	rs296366	T	CT	+/-
16	SULT2A1	rs4149448	G	—	NG
17	SULT2A1	rs4149449	T	CT	+/-
18	SULT2A1	rs4149452	T	CT	+/-
19	SULT2A1	rs8113396	G	—	NG

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report

# Uri rahamim / F

## THYROID

	Gene	rsID	Minor Allele	Genotype	Phenotype
1	CTLA4	rs231775	G	AG	+/-
2	FOXE1	rs10984009	A	GG	-/-
3	FOXE1	rs1867277	A	GG	-/-
4	FOXE1	rs7043516	C	—	NG

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**



## Gene Definitions

### RARA [ +/- ]

Gene mutations can be acquired during a person's lifetime and are present only in certain cells. These mutations are called somatic mutations, and they are not inherited. A somatic mutation involving the RARA gene causes acute promyelocytic leukemia, a cancer of the blood forming tissue (bone marrow). Acute promyelocytic leukemia is characterized by an accumulation of promyelocytes in the bone marrow. A rearrangement (translocation) of genetic material between chromosomes 15 and 17, written as t(15;17), fuses part of the RARA gene on chromosome 17 with part of another gene on chromosome 15 called PML . The protein produced from this fused gene, the PML-RAR\u03b1 protein, functions differently than the protein products of the normal PML and RARA genes. The PML-RAR\u03b1 protein binds to DNA and represses gene transcription, like the normal RAR\u03b1 protein. However, the PML-RAR\u03b1 protein does not respond to the signal to induce transcription of genes, so the genes remain repressed. Additionally, the function of the PML protein, the product of the PML gene, is disrupted. The PML protein blocks cell growth and division (proliferation) and induces self-destruction (apoptosis) in combination with other proteins. However, the PML-RAR\u03b1 protein does not block proliferation or induce apoptosis. The PML-RAR\u03b1 protein blocks the differentiation of blood cells at the promyelocyte stage and allows abnormal cell proliferation. As a result, excess promyelocytes accumulate in the bone marrow and normal white blood cells cannot form, leading to acute promyelocytic leukemia.

### CETP [ +/- ]

The protein encoded by this gene is found in plasma, where it is involved in the transfer of cholesteryl ester from high density lipoprotein (HDL) to other lipoproteins. Defects in this gene are a cause of hyperalphalipoproteinemia 1 (HALP1). Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2013]

### CYP4V2 [ +/- ]

At least 42 CYP4V2 gene mutations have been identified in people with Bietti crystalline dystrophy, a disorder in which numerous small, yellow or white crystal-like deposits of fatty (lipid) compounds accumulate in the light-sensitive tissue that lines the back of the eye (the retina). The deposits damage the retina, resulting in progressive vision loss. CYP4V2 gene mutations that cause Bietti crystalline dystrophy are predicted to change the structure of the CYP4V2 enzyme in a way that reduces or eliminates its activity. The mutations likely affect lipid breakdown; however, it is unknown how they lead to the specific signs and symptoms of Bietti crystalline dystrophy. For unknown reasons, the severity of the signs and symptoms differs significantly among individuals with the same CYP4V2 gene mutation.

### GP6 [ +/- ]

Normally, if you get hurt, your body forms a blood clot to stop the bleeding. For blood to clot, your body needs cells called platelets and proteins known as clotting factors. If you have a bleeding disorder, you either do not have enough platelets or clotting factors or they don't work the way they should. Bleeding disorders can be the result of other diseases, such as severe liver disease. They can also be inherited. Hemophilia is an inherited bleeding disorder. Bleeding disorders can also be a side effect of medicines.

### HRG [ +/- ]

The protein encoded by this gene was originally identified as a 44-kD glycoprotein that interacts with the NEU/ERBB2 receptor tyrosine kinase to increase its phosphorylation on tyrosine residues. This protein is a signaling protein that

mediates cell-cell interactions and plays critical roles in the growth and development of multiple organ systems. It is known that an extraordinary variety of different isoforms are produced from this gene through alternative promoter usage and splicing. These isoforms are tissue-specifically expressed and differ significantly in their structure, and thereby these isoforms are classified into types I, II, III, IV, V and VI. The gene dysregulation has been linked to diseases such as cancer, schizophrenia and bipolar disorder (BPD). [provided by RefSeq, May 2009]

### ITGB3 [ +/- ]

Platelets are little pieces of blood cells. Platelets help wounds heal and prevent bleeding by forming blood clots. Your bone marrow makes platelets. Problems can result from having too few or too many platelets, or from platelets that do not work properly. If your blood has a low number of platelets, you can be at risk for mild to serious bleeding. If your blood has too many platelets, you may have a higher risk of blood clots. With other platelet disorders, the platelets do not work as they should. For example, in von Willebrand Disease, the platelets cannot stick together or cannot attach to blood vessel walls. This can cause excessive bleeding. Treatment of platelet disorders depends on the cause. NIH: National Heart, Lung, and Blood Institute

### NR1I2 [ +/- ]

This gene product belongs to the nuclear receptor superfamily, members of which are transcription factors characterized by a ligand-binding domain and a DNA-binding domain. The encoded protein is a transcriptional regulator of the cytochrome P450 gene CYP3A4, binding to the response element of the CYP3A4 promoter as a heterodimer with the 9-cis retinoic acid receptor RXR.

### CTH [ +/- ]

This gene encodes a cytoplasmic enzyme in the trans-sulfuration pathway that converts cystathione derived from methionine into cysteine. Glutathione synthesis in the liver is dependent upon the availability of cysteine. Mutations in this gene cause cystathioninuria. Alternative splicing of this gene results in three transcript variants encoding different isoforms. [provided by RefSeq, Jun 2010]

### CYP1B1 [ +/- ]

This gene encodes an estrogen receptor, a ligand-activated transcription factor composed of several domains important for hormone binding, DNA binding, and activation of transcription. The protein localizes to the nucleus where it may form a homodimer or a heterodimer with estrogen receptor 2. Estrogen and its receptors are essential for sexual development and reproductive function, but also play a role in other tissues such as bone. Estrogen receptors are also involved in pathological processes including breast cancer, endometrial cancer, and osteoporosis. Alternative promoter usage and alternative splicing result in dozens of transcript variants, but the full-length nature of many of these variants has not been determined. [provided by RefSeq, Mar 2014]

### CYP2C9 [ +/- ]

Antidepressants are medicines that treat depression. Your doctor can prescribe them for you. They work to balance some of the natural chemicals in our brains. It may take several weeks for them to help. There are several types of antidepressants. You and your doctor may have to try a few before finding what works best for you. Antidepressants may cause mild side effects that usually do not last long. These may include headache, nausea, sleep problems, restlessness, and sexual problems. Tell your doctor if you have any side effects. You should also let your doctor know if you take any other medicines, vitamins, or herbal supplements. It is important to keep taking your medicines, even if you feel better. Do not stop taking your medicines without talking to your doctor. You often need to stop antidepressants

## CYP2D6 [ +/- ]

Caffeine is a bitter substance found in coffee, tea, soft drinks, chocolate, kola nuts, and certain medicines. It has many effects on the body's metabolism, including stimulating the central nervous system. This can make you more alert and give you a boost of energy. For most people, the amount of caffeine in two to four cups of coffee a day is not harmful. However, too much caffeine can cause problems. It canMake you jittery and shakyMake it hard to fall asleep or stay asleepCause headaches or dizzinessMake your heart beat faster or cause abnormal heart rhythmsCause dehydrationMake you dependent on it so you need to take more of it. If you stop using caffeine, you could get withdrawal symptoms.Some people are more sensitive to the effects of caffeine than others. They should limit their use of caffeine. So should pregnant and nursing women. Certain drugs and supplements may interact with caffeine. If you have questions about whether caffeine is safe for you, talk with your health care provider.Food and Drug Administration

## NAT2 [ +/- ]

Note: Loci in other organisms that are functionally homologous to this one are validly referred to as both NAT1 and NAT2; i.e., the functional homologs of NAT1 include mouse and rat Nat2, while the functional homologs of human NAT2 include mouse and rat Nat1. Name:sequence associations are consistent with current use in the field. [27 Apr 2009]

## HLA-DQA1 [ +/- ]

SS2; BTN7; BTL-II; HSBLMHC1

## HLA-DPA1 [ +/- ]

SS2; BTN7; BTL-II; HSBLMHC1

## HLA-DQA2 [ +/+ ]

SS2; BTN7; BTL-II; HSBLMHC1

## HLA-DRB1 [ +/- ]

SS2; BTN7; BTL-II; HSBLMHC1

## IFIH1 [ +/+ ]

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein that is upregulated in response to treatment with beta-interferon and a protein kinase C-activating compound, mezerein. Irreversible reprogramming of melanomas can be achieved by treatment with both these agents; treatment with either agent alone only achieves reversible differentiation. Genetic variation in this gene is associated with diabetes mellitus insulin-dependent type 19. [provided

### IRF5 [ +/- ]

Studies have associated normal variations in the IRF5 gene with an increased risk of several autoimmune disorders. Autoimmune disorders occur when the immune system malfunctions and attacks the body's tissues and organs. These disorders include systemic lupus erythematosus, Sjögren syndrome, and rheumatoid arthritis. There is some evidence that certain variations of the IRF5 gene are associated with increased activity of the gene and elevated cytokines. However, it is unknown what role, if any, these effects play in the increased risk of autoimmune disorders. Researchers believe that a combination of genetic and environmental factors may contribute to the development of these conditions.

### MTC03P1 [ +/+ ]

MTC03P1

### PSMB8 [ +/- ]

At least one mutation in the PSMB8 gene has been found to cause Nakajo-Nishimura syndrome, a condition that has been described only in the Japanese population. The identified mutation changes a single protein building block (amino acid) in the protein produced from the PSMB8 gene, replacing the amino acid glycine with the amino acid valine at protein position 201 (written as Gly201Val or G201V). This mutation greatly reduces the production of this protein, which impairs the normal assembly of immunoproteasomes and causes the immune system to malfunction. For unknown reasons, the malfunctioning immune system triggers abnormal inflammation that can damage tissues throughout the body. Abnormal inflammation likely underlies many of the signs and symptoms of Nakajo-Nishimura syndrome, including the development of red, swollen lumps (nodular erythema) on the skin, recurrent fevers, joint problems, and an enlarged liver and spleen (hepatosplenomegaly). It is less clear how mutations in the PSMB8 gene lead to other characteristic features of the condition, including muscle weakness and wasting and a loss of fatty tissue (lipodystrophy), mainly in the upper body. Because the protein produced from the PSMB8 gene may be involved in the maturation of adipocytes, studies suggest that a shortage of this protein may interfere with the normal development and function of these cells.

### C3 [ +/- ]

At least one mutation in the C3 gene has been found to cause dense deposit disease. This condition, which was formerly known as membranoproliferative glomerulonephritis type II, is a form of progressive kidney (renal) disease. The identified mutation deletes two amino acids from the C3 protein. This genetic change is described as a "gain-of-function" mutation because it leads to abnormal activation of the complement system. The overactive system creates debris that builds up in and damages certain structures in the kidneys. These structures, called glomeruli, are clusters of tiny blood vessels that help filter waste products from the blood. Damage to glomeruli prevents the kidneys from filtering waste products normally and can lead to end-stage renal disease (ESRD), a life-threatening failure of kidney function. Several normal variants (polymorphisms) in the C3 gene have also been associated with an increased risk of developing dense deposit disease. In particular, the C3F allotype is seen more frequently in people with this condition than in the general population. Researchers are working to determine how the C3F allotype influences disease risk.

### FCER1A [ +/+ ]

The immunoglobulin epsilon receptor (IgE receptor) is the initiator of the allergic response. When two or more high-affinity IgE receptors are brought together by allergen-bound IgE molecules, mediators such as histamine that are responsible for allergy symptoms are released. This receptor is comprised of an alpha subunit, a beta subunit, and two gamma subunits. The protein encoded by this gene represents the alpha subunit. [provided by RefSeq, Aug 2011]

### IL5 [ +/- ]

This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation, maturation, recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor, which is a heterodimer, whose beta subunit is shared with the receptors for interleukin 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4), interleukin 13 (IL13), and CSF2. This gene, IL4, and IL13 may be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31. [provided by RefSeq, Jul 2013]

### RAG1 [ +/+ ]

Your immune system is a complex network of cells, tissues, and organs that work together to defend against germs. It helps your body to recognize these "foreign" invaders. Then its job is to keep them out, or if it can't, to find and destroy them. If your immune system cannot do its job, the results can be serious. Disorders of the immune system include Allergy and asthma - immune responses to substances that are usually not harmful Immune deficiency diseases - disorders in which the immune system is missing one or more of its parts Autoimmune diseases - diseases causing your immune system to attack your own body's cells and tissues by mistake NIH: National Institute of Allergy and Infectious Diseases

### SOCS1 [ +/- ]

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons (PMIDs: 12032546, 20937277). [provided by RefSeq, Feb 2013]

### FCGR2A [ +/- ]

Malaria is a serious disease caused by a parasite. You get it when an infected mosquito bites you. Malaria is a major cause of death worldwide, but it is almost wiped out in the United States. The disease is mostly a problem in developing countries with warm climates. If you travel to these countries, you are at risk. There are four different types of malaria caused by four related parasites. The most deadly type occurs in Africa south of the Sahara Desert. Malaria symptoms include chills, flu-like symptoms, fever, vomiting, diarrhea, and jaundice. A blood test can diagnose it. It can be life-threatening. However, you can treat malaria with drugs. The type of drug depends on which kind of malaria you have and where you were infected. Malaria can be prevented. When traveling to malaria-prone regions See your doctor for medicines that protect you Wear insect repellent with DEET Cover up Sleep under mosquito netting Centers for Disease Control and Prevention

### GSTM3 [ +/- ]

Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase that belongs to the mu class. The mu class of enzymes functions in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs. Mutations of this class mu gene have been linked with a slight increase in a number of cancers, likely due to exposure with environmental toxins. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2008]

### ACE [ +/+ ]

This gene encodes an enzyme involved in catalyzing the conversion of angiotensin I into a physiologically active peptide angiotensin II. Angiotensin II is a potent vasopressor and aldosterone-stimulating peptide that controls blood pressure and fluid-electrolyte balance. This enzyme plays a key role in the renin-angiotensin system. Many studies have associated the presence or absence of a 287 bp Alu repeat element in this gene with the levels of circulating enzyme or cardiovascular pathophysiology. Multiple alternatively spliced transcript variants encoding different isoforms have been identified, and two most abundant spliced variants encode the somatic form and the testicular form, respectively, that are equally active. [provided by RefSeq, May 2010]

### AGT [ +/+ ]

At least six mutations in the AGT gene have been found to cause a severe kidney disorder called renal tubular dysgenesis. This condition is characterized by abnormal kidney development before birth, the inability to produce urine (anuria), and severe low blood pressure (hypotension). These problems result in a reduction of amniotic fluid (oligohydramnios), which leads to a set of birth defects known as the Potter sequence. Renal tubular dysgenesis can be caused by mutations in both copies of any of the genes involved in the renin-angiotensin system. Most of the mutations in the AGT gene that cause this disorder change single protein building blocks (amino acids) in the angiotensinogen protein. These changes occur in a region of the protein that is necessary for its conversion to angiotensin I. It is thought that the altered angiotensinogen cannot be converted, leading to a nonfunctional renin-angiotensin system. Without this system, the kidneys cannot control blood pressure. Because of low blood pressure, the flow of blood is reduced (hypoperfusion), and the body does not get enough oxygen during fetal development. As a result, kidney development is impaired, leading to the features of renal tubular dysgenesis.

### BHMT [ +/- ]

This gene encodes a cytosolic enzyme that catalyzes the conversion of betaine and homocysteine to dimethylglycine and methionine, respectively. Defects in this gene could lead to hyperhomocyst(e)inemia, but such a defect has not yet been observed. [provided by RefSeq, Jul 2008]

### C1orf167 [ +/- ]

C1ORF167

### CBS A360A [ +/- ]

More than 150 mutations that cause homocystinuria have been identified in the CBS gene. Most of these mutations change single amino acids in cystathionine beta-synthase. The most common mutation substitutes the amino acid threonine for the amino acid isoleucine at position 278 in the enzyme (written as Ile278Thr or I278T). Another common mutation, which is the most frequent cause of homocystinuria in the Irish population, replaces the amino acid glycine with the amino acid serine at position 307 (written as Gly307Ser or G307S). These mutations disrupt the normal function of cystathionine beta-synthase. As a result, homocysteine and other potentially toxic compounds build up in the blood, and homocysteine is excreted in urine. Researchers have not determined how excess homocysteine leads to the signs and symptoms of homocystinuria.

### COMT [ +/- ]

The characteristic signs and symptoms of 22q11.2 deletion syndrome result from a deletion of a small piece of chromosome 22. The chromosomal region that is typically deleted contains 30 to 40 genes, including the COMT gene. As a result of the deletion, people with this disorder have only one copy of the COMT gene in each cell instead of the usual two copies. A loss of one copy of the COMT gene in each cell leads to abnormal regulation of catechol-O-methyltransferase levels in the brain. Researchers believe that changes involving this enzyme in the prefrontal cortex may help explain the increased risk of behavioral problems and mental illness associated with 22q11.2 deletion syndrome. Little is known, however, about the relationship between catechol-O-methyltransferase activity and the specific mental and emotional problems characteristic of this condition. People with 22q11.2 deletion syndrome are much more likely than people without the condition to develop schizophrenia, depression, anxiety, and bipolar disorder.

### COMT H62H [ +/- ]

The characteristic signs and symptoms of 22q11.2 deletion syndrome result from a deletion of a small piece of chromosome 22. The chromosomal region that is typically deleted contains 30 to 40 genes, including the COMT gene. As a result of the deletion, people with this disorder have only one copy of the COMT gene in each cell instead of the usual two copies. A loss of one copy of the COMT gene in each cell leads to abnormal regulation of catechol-O-methyltransferase levels in the brain. Researchers believe that changes involving this enzyme in the prefrontal cortex may help explain the increased risk of behavioral problems and mental illness associated with 22q11.2 deletion syndrome. Little is known, however, about the relationship between catechol-O-methyltransferase activity and the specific mental and emotional problems characteristic of this condition. People with 22q11.2 deletion syndrome are much more likely than people without the condition to develop schizophrenia, depression, anxiety, and bipolar disorder.

### COMT V158M [ +/- ]

The characteristic signs and symptoms of 22q11.2 deletion syndrome result from a deletion of a small piece of chromosome 22. The chromosomal region that is typically deleted contains 30 to 40 genes, including the COMT gene. As a result of the deletion, people with this disorder have only one copy of the COMT gene in each cell instead of the usual two copies. A loss of one copy of the COMT gene in each cell leads to abnormal regulation of catechol-O-methyltransferase levels in the brain. Researchers believe that changes involving this enzyme in the prefrontal cortex may help explain the increased risk of behavioral problems and mental illness associated with 22q11.2 deletion syndrome. Little is known, however, about the relationship between catechol-O-methyltransferase activity and the specific mental and emotional problems characteristic of this condition. People with 22q11.2 deletion syndrome are much more likely than people without the condition to develop schizophrenia, depression, anxiety, and bipolar disorder.

### DAO [ +/- ]

Schizophrenia is a serious brain illness. People who have it may hear voices that aren't there. They may think other people are trying to hurt them. Sometimes they don't make sense when they talk. The disorder makes it hard for them to keep a job or take care of themselves. Symptoms of schizophrenia usually start between ages 16 and 30. Men often develop symptoms at a younger age than women. People usually do not get schizophrenia after age 45. There are

three types of symptoms: Psychotic symptoms distort a person's thinking. These include hallucinations (hearing or seeing things that are not there), delusions (beliefs that are not true), trouble organizing thoughts, and strange movements. "Negative" symptoms make it difficult to show emotions and to function normally. A person may seem depressed and withdrawn. Cognitive symptoms affect the thought process. These include trouble using information, making decisions, and paying attention. No one is sure what causes schizophrenia. Your genes, environment, and brain chemistry may play a role. There is no cure. Medicine can help control many of the symptoms. You may need to try different medicines to see which works best. You should stay on your medicine for as long as your doctor recommends. Additional treatments can help you deal with your illness from day to day. These include therapy, family education, rehabilitation, and skills training. NIH: National Institute of Mental Health

### DHFR [ +/+ ]

Dihydrofolate reductase converts dihydrofolate into tetrahydrofolate, a methyl group shuttle required for the de novo synthesis of purines, thymidylic acid, and certain amino acids. While the functional dihydrofolate reductase gene has been mapped to chromosome 5, multiple intronless processed pseudogenes or dihydrofolate reductase-like genes have been identified on separate chromosomes. Dihydrofolate reductase deficiency has been linked to megaloblastic anemia. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2014]

### FUT2 [ +/- ]

The protein encoded by this gene is a Golgi stack membrane protein that is involved in the creation of a precursor of the H antigen, which is required for the final step in the soluble A and B antigen synthesis pathway. This gene is one of two encoding the galactoside 2-L-fucosyltransferase enzyme. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008]

### GAD1 [ +/- ]

Cerebral palsy is a group of disorders that affect a person's ability to move and to maintain balance and posture. The disorders appear in the first few years of life. Usually they do not get worse over time. People with cerebral palsy may have difficulty walking. They may also have trouble with tasks such as writing or using scissors. Some have other medical conditions, including seizure disorders or mental impairment. Cerebral palsy happens when the areas of the brain that control movement and posture do not develop correctly or get damaged. Early signs of cerebral palsy usually appear before 3 years of age. Babies with cerebral palsy are often slow to roll over, sit, crawl, smile, or walk. Some babies are born with cerebral palsy; others get it after they are born. There is no cure for cerebral palsy, but treatment can improve the lives of those who have it. Treatment includes medicines, braces, and physical, occupational and speech therapy. NIH: National Institute of Neurological Disorders and Stroke

### GIF [ +/- ]

This gene encodes a lymphokine involved in cell-mediated immunity, immunoregulation, and inflammation. It plays a role in the regulation of macrophage function in host defense through the suppression of anti-inflammatory effects of glucocorticoids. This lymphokine and the JAB1 protein form a complex in the cytosol near the peripheral plasma membrane, which may indicate an additional role in integrin signaling pathways. [provided by RefSeq, Jul 2008]

### MTHFD1 [ +/- ]

Neural tube defects are birth defects of the brain, spine, or spinal cord. They happen in the first month of pregnancy, often before a woman even knows that she is pregnant. The two most common neural tube defects are spina bifida and

anencephaly. In spina bifida, the fetal spinal column doesn't close completely. There is usually nerve damage that causes at least some paralysis of the legs. In anencephaly, most of the brain and skull do not develop. Babies with anencephaly are either stillborn or die shortly after birth. Another type of defect, Chiari malformation, causes the brain tissue to extend into the spinal canal. The exact causes of neural tube defects aren't known. You're at greater risk of having an infant with a neural tube defect if you are obese, have poorly controlled diabetes, take certain antiseizure medicines, or don't get enough folic acid, a type of B vitamin, before and during pregnancy. Getting enough folic acid prevents most neural tube defects. Neural tube defects are usually diagnosed before the infant is born, through lab or imaging tests. There is no cure for neural tube defects. The nerve damage and loss of function that are present at birth are usually permanent. However, a variety of treatments can sometimes prevent further damage and help with complications. NIH: National Institute of Child Health and Human Development

### MTHFD1L [ +/- ]

The protein encoded by this gene is involved in the synthesis of tetrahydrofolate (THF) in the mitochondrion. THF is important in the de novo synthesis of purines and thymidylate and in the regeneration of methionine from homocysteine. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2011]

### MTHFR [ +/- ]

At least 40 mutations in the MTHFR gene have been identified in people with homocystinuria, a disorder in which the body is unable to process certain amino acids properly. Most of these mutations change single amino acids in methylenetetrahydrofolate reductase. These changes impair the function of the enzyme, and some cause the enzyme to be turned off (inactivated). Other mutations lead to the production of an abnormally small, nonfunctional version of the enzyme. Without functional methylenetetrahydrofolate reductase, homocysteine cannot be converted to methionine. As a result, homocysteine builds up in the bloodstream, and the amount of methionine is reduced. Some of the excess homocysteine is excreted in urine. Researchers have not determined how altered levels of homocysteine and methionine lead to the various health problems affecting multiple parts of the body in people with homocystinuria.

### MTHFS [ +/- ]

The protein encoded by this gene is an enzyme that catalyzes the conversion of 5-formyltetrahydrofolate to 5,10-methylenetetrahydrofolate, a precursor of reduced folates involved in 1-carbon metabolism. An increased activity of the encoded protein can result in an increased folate turnover rate and folate depletion. Three transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jun 2011]

### MTR A2756G [ +/- ]

More than 20 mutations in the MTR gene have been identified in people with homocystinuria. Many of these mutations lead to the production of an abnormally small, nonfunctional version of methionine synthase. Other mutations change single amino acids in the enzyme. One of the most common mutations replaces the amino acid proline with the amino acid leucine at position 1173 (written as Pro1173Leu or P1173L), resulting in an enzyme with reduced function. Without functional methionine synthase, homocysteine cannot be converted to methionine. As a result, homocysteine builds up in the bloodstream, and the amount of methionine is reduced. Some of the excess homocysteine is excreted in urine. Researchers have not determined how altered levels of homocysteine and methionine lead to the health problems associated with homocystinuria.

### MTRR A664A [ +/- ]

At least 20 mutations in the MTRR gene have been identified in people with homocystinuria. Some of these mutations change single amino acids in methionine synthase reductase. Other mutations lead to an abnormally small, nonfunctional version of the enzyme. All these mutations prevent the enzyme from functioning normally. Without methionine synthase reductase, methionine synthase cannot convert homocysteine to methionine. As a result, homocysteine builds up in the bloodstream, and the amount of methionine is reduced. Some of the excess homocysteine is excreted in urine. Researchers have not determined how altered levels of homocysteine and methionine lead to the health problems associated with homocystinuria.

### MTRR K350A [ +/- ]

At least 20 mutations in the MTRR gene have been identified in people with homocystinuria. Some of these mutations change single amino acids in methionine synthase reductase. Other mutations lead to an abnormally small, nonfunctional version of the enzyme. All these mutations prevent the enzyme from functioning normally. Without methionine synthase reductase, methionine synthase cannot convert homocysteine to methionine. As a result, homocysteine builds up in the bloodstream, and the amount of methionine is reduced. Some of the excess homocysteine is excreted in urine. Researchers have not determined how altered levels of homocysteine and methionine lead to the health problems associated with homocystinuria.

### NOS2 [ +/- ]

Blood pressure is the force of your blood pushing against the walls of your arteries. Each time your heart beats, it pumps blood into the arteries. Your blood pressure is highest when your heart beats, pumping the blood. This is called systolic pressure. When your heart is at rest, between beats, your blood pressure falls. This is called diastolic pressure. Your blood pressure reading uses these two numbers. Usually the systolic number comes before or above the diastolic number. A reading of 119/79 or lower is normal blood pressure. 140/90 or higher is high blood pressure. Between 120 and 139 for the top number, or between 80 and 89 for the bottom number is called prehypertension. Prehypertension means you may end up with high blood pressure, unless you take steps to prevent it. High blood pressure usually has no symptoms, but it can cause serious problems such as stroke, heart failure, heart attack and kidney failure. You can control high blood pressure through healthy lifestyle habits and taking medicines, if needed. NIH: National Heart, Lung, and Blood Institute

### NOS3 [ +/- ]

Alzheimer's disease (AD) is the most common form of dementia among older people. Dementia is a brain disorder that seriously affects a person's ability to carry out daily activities. AD begins slowly. It first involves the parts of the brain that control thought, memory and language. People with AD may have trouble remembering things that happened recently or names of people they know. A related problem, mild cognitive impairment (MCI), causes more memory problems than normal for people of the same age. Many, but not all, people with MCI will develop AD. In AD, over time, symptoms get worse. People may not recognize family members or have trouble speaking, reading or writing. They may forget how to brush their teeth or comb their hair. Later on, they may become anxious or aggressive, or wander away from home. Eventually, they need total care. This can cause great stress for family members who must care for them. AD usually begins after age 60. The risk goes up as you get older. Your risk is also higher if a family member has had the disease. No treatment can stop the disease. However, some drugs may help keep symptoms from getting worse for a limited time. NIH: National Institute on Aging

### PEMT [ +/+ ]

This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast, stomach and pancreas. This protein is proteolytically cleaved

into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Feb 2011]

## SOD2 [ +/+ ]

This gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of this cytokine. [provided by RefSeq, Jul 2008]

## SOD3 [ +/- ]

This gene encodes an extracellular matrix glycoprotein that is characterized by an N-terminal microfibril interface domain, a coiled-coiled alpha-helical domain, a collagenous domain and a C-terminal globular C1q domain. The encoded protein associates with elastic fibers at the interface between elastin and microfibrils and may play a role in the development of elastic tissues including large blood vessels, dermis, heart and lung. [provided by RefSeq, Sep 2009]

## TCN1 [ +/- ]

This gene encodes a member of the vitamin B12-binding protein family. This family of proteins, alternatively referred to as R binders, is expressed in various tissues and secretions. This protein is a major constituent of secondary granules in neutrophils and facilitates the transport of cobalamin into cells. [provided by RefSeq, Jul 2008]

## TCN2 [ +/- ]

More than 20 mutations in the TCN2 gene have been found to cause transcobalamin deficiency. This condition impairs the transport of cobalamin from the bloodstream to cells throughout the body. Affected individuals have difficulty gaining weight and growing at the expected rate (failure to thrive), vomiting, diarrhea, a shortage of all types of blood cells, and neurological problems. Many TCN2 gene mutations lead to a complete or near-complete lack (deficiency) of transcobalamin. Other TCN2 gene mutations result in a transcobalamin protein that cannot bind to cobalamin or a protein that cannot bind to the receptor at the surface of cells. The resulting lack of cobalamin within cells interferes with the functioning of certain enzymes, which impacts many cell activities. As a result, a wide range of signs and symptoms characteristic of transcobalamin deficiency can develop.

## NDUFS7 [ +/- ]

A genetic brain disorder is caused by a variation or a mutation in a gene. A variation is a different form of a gene. A mutation is a change in a gene. Genetic brain disorders affect the development and function of the brain. Some genetic brain disorders are due to random gene mutations or mutations caused by environmental exposure, such as cigarette smoke. Other disorders are inherited, which means that a mutated gene or group of genes is passed down through a family. They can also be due to a combination of both genetic changes and other outside factors. Some examples of genetic brain disorders include Leukodystrophies, Phenylketonuria, Tay-Sachs disease, Wilson disease. Many people with genetic brain disorders fail to produce enough of certain proteins that influence brain development and function. These brain disorders can cause serious problems that affect the nervous system. Some have treatments to control

symptoms. Some are life-threatening.

### NDUFS8 [ +/+ ]

A genetic brain disorder is caused by a variation or a mutation in a gene. A variation is a different form of a gene. A mutation is a change in a gene. Genetic brain disorders affect the development and function of the brain. Some genetic brain disorders are due to random gene mutations or mutations caused by environmental exposure, such as cigarette smoke. Other disorders are inherited, which means that a mutated gene or group of genes is passed down through a family. They can also be due to a combination of both genetic changes and other outside factors. Some examples of genetic brain disorders include Leukodystrophies, Phenylketonuria, Tay-Sachs disease, and Wilson disease. Many people with genetic brain disorders fail to produce enough of certain proteins that influence brain development and function. These brain disorders can cause serious problems that affect the nervous system. Some have treatments to control symptoms. Some are life-threatening.

### ATG16L1 [ +/+ ]

At least one variation in the ATG16L1 gene is associated with an increased risk of Crohn disease, particularly a form of the disorder that affects the lower part of the small intestine (the ileum). This increased risk has been found primarily in Caucasian (white) populations. The identified ATG16L1 variation changes a single protein building block (amino acid) in a critical region of the autophagy related 16-like 1 protein. Specifically, it replaces the amino acid threonine with the amino acid alanine at protein position 300 (written as Thr300Ala or T300A). The effects of variations in the ATG16L1 gene on Crohn disease risk are unclear. Changes in this gene may affect the autophagy process, allowing worn-out cell parts and harmful bacteria to persist when they would otherwise be destroyed. These cell components and bacteria may trigger an inappropriate immune system response, leading to chronic inflammation in the intestinal walls and the digestive problems characteristic of Crohn disease. Researchers continue to study the relationship between changes in the ATG16L1 gene and a person's risk of developing this disorder.

### IL13 [ +/- ]

Each spring, summer, and fall, trees, weeds, and grasses release tiny pollen grains into the air. Some of the pollen ends up in your nose and throat. This can trigger a type of allergy called hay fever. Symptoms can include sneezing, often with a runny or clogged nose, coughing and postnasal drip, itchy eyes, nose and throat, red and watery eyes, dark circles under the eyes. Your health care provider may diagnose hay fever based on a physical exam and your symptoms. Sometimes skin or blood tests are used. Taking medicines and using nasal sprays can relieve symptoms. You can also rinse out your nose, but be sure to use distilled or sterilized water with saline. Allergy shots can help make you less sensitive to pollen and provide long-term relief. NIH: National Institute of Allergy and Infectious Diseases

### IL4R [ +/- ]

In the early 1980s, when the HIV/AIDS epidemic began, patients rarely lived longer than a few years. But today, there are many effective medicines to fight the infection, and people with HIV have longer, healthier lives. There are five major types of medicines: Reverse transcriptase (RT) inhibitors - interfere with a critical step during the HIV life cycle and keep the virus from making copies of itself; Protease inhibitors - interfere with a protein that HIV uses to make infectious viral particles; Fusion inhibitors - block the virus from entering the body's cells; Integrase inhibitors - block an enzyme HIV needs to make copies of itself; Multidrug combinations - combine two or more different types of drugs into one. These medicines help people with HIV, but they are not perfect. They do not cure HIV/AIDS. People with HIV infection still have the virus in their bodies. They can still spread HIV to others through unprotected sex and needle sharing, even when they are taking their medicines. NIH: National Institute of Allergy and Infectious Diseases

## SULT2A1 [ +/- ]

This gene encodes a member of the sulfotransferase family. Sulfotransferases aid in the metabolism of drugs and endogenous compounds by converting these substances into more hydrophilic water-soluble sulfate conjugates that can be easily excreted. This protein catalyzes the sulfation of steroids and bile acids in the liver and adrenal glands, and may have a role in the inherited adrenal androgen excess in women with polycystic ovary syndrome. [provided by RefSeq, Mar 2010]

## CTLA4 [ +/- ]

Diabetes is a disease in which your blood glucose, or blood sugar, levels are too high. Glucose comes from the foods you eat. Insulin is a hormone that helps the glucose get into your cells to give them energy. With type 1 diabetes, your body does not make insulin. With type 2 diabetes, the more common type, your body does not make or use insulin well. Without enough insulin, the glucose stays in your blood. You can also have prediabetes. This means that your blood sugar is higher than normal but not high enough to be called diabetes. Having prediabetes puts you at a higher risk of getting type 2 diabetes. Over time, having too much glucose in your blood can cause serious problems. It can damage your eyes, kidneys, and nerves. Diabetes can also cause heart disease, stroke and even the need to remove a limb. Pregnant women can also get diabetes, called gestational diabetes. A blood test can show if you have diabetes. Exercise, weight control and sticking to your meal plan can help control your diabetes. You should also monitor your glucose level and take medicine if prescribed. NIH: National Institute of Diabetes and Digestive and Kidney Diseases

A gene variance report is a graphical representation of genetic raw data, displayed as a color coded chart. Phenotypes are determined based on the presence or absence of variant alleles in your genotype. Alleles are considered variant if they are the minor allele i.e they occur with less frequency (MAF) in the default global population. Having "no variant" alleles (green) is not necessarily "normal" or protective, and having a homozygous phenotype (red) is not always "abnormal". What is a normal or abnormal phenotype should NOT be determined solely based on this variance report. The significance of your phenotypes should be assessed by reviewing related genome wide studies for context and in consultation with a qualified health practitioner or genetics specialist. All alleles are reported in reference to the forward strand. rsIDs and genotype information are obtained from the genetic raw data prepared by your personal genomic service. Minor allele frequency (MAF), RefSNP and gene variation/SNP names are obtained directly from dbSNP which is a free public archive for genetic variation maintained by the NCBI <http://www.ncbi.nlm.nih.gov/snp/>.

Disclaimer: (1) No advice. This website contains general information about medical conditions and treatments. The information is not advice, and should not be treated as such. (2) No warranties The information on this website is provided without any representations or warranties, express or implied. We make no representations warranties in relation to the information on this website. Without prejudice to the generality of the foregoing paragraph, we do not warrant that: (a) the information on this website will be constantly available, or available at all; or (b) the information on this website is complete, true, up-to date. (3) Professional assistance You must not rely on the information on this website as an alternative to medical advice from your doctor or other professional healthcare provider. If you have any specific questions about any medical matter you should consult your doctor or other professional healthcare provider. If you think you may be suffering from any medical condition you should seek immediate medical attention. You should never delay seeking medical advice, disregard medical advice, or discontinue medical treatment because of information on this website. (4) Limiting our liability. Except as provided by applicable law, nothing in this disclaimer will render us liable for death or personal injury resulting from review or use of information available on this website. (a) limit or exclude our liability for death or personal injury resulting from negligence; . Because of the potential harms from genetic test information and the need for patients and their families to receive appropriate information for decision making, patients with possible inherited diseases should undergo genetic testing only in the context of genetic counseling.

Report generated by Livewello Gene Variance Software. <http://livewello.com/genetics> v2.7

**Do not make any decisions about your health solely based on the information contained in this report. Always consult with a licensed and experienced health practitioner when you receive your report**

Compiled on Tue May 12 15:25:10 2015